

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP04/014170

International filing date: 13 December 2004 (13.12.2004)

Document type: Certified copy of priority document

Document details: Country/Office: EP
Number: 04013328.2
Filing date: 05 June 2004 (05.06.2004)

Date of receipt at the International Bureau: 28 January 2005 (28.01.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

04013328.2

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 04013328.2
Demande no:

Anmeldetag:
Date of filing: 05.06.04
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Epigenomics AG
Kastanienallee 24
10435 Berlin
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

PITX2 - a marker to predict survival of patients diagnosed with breast cell
proliferative disease

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

G12Q1/68

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PL PT RO SE SI SK TR LI

PITX2 - a marker to predict survival of patients diagnosed with breast cell proliferative disease

The present invention relates to methods for predicting the survival of a human being diagnosed with a cell proliferative disorder of the breast tissues, characterized by a step of determining the expression level of PITX2 or the genetic or the epigenetic modifications of the genomic DNA associated with the gene PITX2. The invention also relates to sequences, oligonucleotides and antibodies which can be used within the described methods.

Field of the Invention

BREAST CANCER SURVIVAL

In European and American women, breast cancer is the most frequently diagnosed cancer and the second leading cause of cancer death. In women aged 40-55, breast cancer is the leading cause of death (Greenlee *et al.*, 2000). In 2002, there were 204,000 new cases of breast cancer in the US and a comparable number in Europe.

Breast cancer is defined as the uncontrolled proliferation of cells within breast tissues. Breasts are comprised of 15 to 20 lobes joined together by ducts. Cancer arises most commonly in the duct, but is also found in the lobes with the rarest type of cancer termed inflammatory breast cancer. It will be appreciated by those skilled in the art that there exists a continuing need to improve methods of early detection, classification and treatment of breast cancers. In contrast to the detection of some other common cancers such as cervical and dermal there are inherent difficulties in classifying and detecting breast cancers.

The first step of any treatment is the assessment of the patient's condition comparative to defined classifications of the disease. However the value of such a system is inherently dependent upon the quality of the classification. Breast cancers are staged according to their size, location and occurrence of metastasis. Methods of treatment include the use of surgery, radiation therapy, chemotherapy and endocrine therapy, which are also used as adjuvant therapies to surgery. In general, more aggressive disease should be treated with more aggressive therapies.

Although the vast majority of early cancers are operable, i.e. the tumor can be completely removed by surgery, about one third of the patients with lymph-node negative diseases and about 50-60% of patients with node-positive disease will develop metastases during follow-up.

Based on this observation, systemic adjuvant treatment has been introduced for both node-positive and node-negative breast cancers. Systemic adjuvant therapy is administered after surgical removal of the tumour, and has been shown to reduce the risk of recurrence significantly. Several types of adjuvant treatment are available: endocrine treatment, also called hormone treatment (for hormone receptor positive tumours), different chemotherapy regimens, and antibody treatments based on novel agents like Herceptin (an antibody to an epidermal growth factor receptor).

The growth of the majority of breast cancers (app. 70-80%) is dependent on the presence of estrogen. Therefore, one important target for adjuvant therapy is the removal of estrogen (e.g. by ovarian ablation) or the blocking of its synthesis or the blocking of its actions on the tumor cells either by blocking the receptor with competing substances (e.g. Tamoxifen) or by inhibiting the conversion of androgen into estrogen (e.g. aromatase inhibitors). This type of treatment is called "endocrine treatment". Endocrine treatment is thought to be efficient only in tumors that express hormone receptors (the estrogen receptor (ER) and/or the progesterone receptor (PR)). Currently, the vast majority of women with hormone receptor positive breast cancer receive some form of endocrine treatment, independent of their nodal status. The most frequently used drug in this scenario is Tamoxifen.

However, even in hormone receptor positive patients, not all patients benefit from endocrine treatment. Adjuvant endocrine therapy reduces mortality rates by 22% while response rates to endocrine treatment in the metastatic (advanced) setting are 50 to 60%.

Since Tamoxifen has relatively few side effects, treatment may be justified even for patients with low likelihood of benefit. However, these patients may require additional, more aggressive adjuvant treatment. Even in earliest and least aggressive tumours, such as node-negative, hormone receptor positive tumours, about 21% of patients relapse within 10 years after initial diagnosis if they receive Tamoxifen monotherapy only, as adjuvant treatment (Lancet. 1998

May 16;351(9114):1451-67. Tamoxifen for early breast cancer: an overview of the randomised trials. Early Breast Cancer Trialists' Collaborative Group.). Similarly, some patients with hormone receptor negative disease may be treated sufficiently with surgery and potentially radiotherapy alone, whereas others may require additional chemotherapy.

Several cytotoxic regimens have shown to be effective in reducing the risk of relapse in breast cancer (Mansour *et al.*, 1998). According to current treatment guidelines, most node-positive patients receive adjuvant chemotherapy both in the US and Europe, since the risk of relapse is considerable. Nevertheless, not all patients do relapse, and there is a proportion of patients who would never have relapsed even without chemotherapy, but who nevertheless receive chemotherapy due to the currently used criteria. In hormone receptor positive patients, chemotherapy is usually given before endocrine treatment, whereas hormone receptor negative patients receive only chemotherapy.

The situation for node-negative patients is particularly complex. In the US, cytotoxic chemotherapy is recommended for node-negative patients, if the tumour is larger than 1 cm. In Europe, chemotherapy is considered for the node-negative cases if one or more risk factors such as tumour size larger than 2 cm, negative hormone receptor status, or tumour grading of three or age <35 is present. In general, there is a tendency to select premenopausal women for additional chemotherapy whereas for postmenopausal women, chemotherapy is often omitted. Compared to endocrine treatment, in particular Tamoxifen or aromatase inhibitors, chemotherapy is highly toxic, with short-term side effects such as nausea, vomiting, bone marrow depression, and long-term effects such as cardiotoxicity and an increased risk for secondary cancers.

It is currently not clear which breast cancer patients should be selected for more aggressive therapy and which would do well without additional aggressive treatment, and clinicians agree that there is a large need for proper selection of patients. The difficulty of selecting the right patients for adjuvant treatment and selecting the right adjuvant treatment, and the lack of suitable criteria is also reflected by a recent study which showed that chemotherapy is used much less frequently than recommended, based on data from the New Mexico Tumor registry (Du *et al.*, 2003). This study provided substantial evidence that there is a need for better selection of patients for chemotherapy or other, more aggressive forms of breast cancer therapy.

This invention is related to a new biomarker, which can be used to solve the problem described above. Based on the observation that methylation of the gene PITX2 (also known as PTX2) in breast tumour tissue, obtained from the surgically removed tumour, or obtained from biopsy material prior to the removal, is correlated with the survival time of breast cancer patients treated with Tamoxifen monotherapy, we invented a tool allowing a better selection of patients for adjuvant therapy, for example for a cytotoxic therapy (chemotherapy) (besides of or instead of an endocrine treatment like treatment with Tamoxifen or aromatase inhibitors) for breast cancer patients.

It can be concluded that the expression levels of the protein or mRNA also correlate with said prognosis. Therefore, the analysis of either the expression levels of PITX2 protein, or PITX2 mRNA or the analysis of the patient's individual genetic or epigenetic modification of the gene PITX2 – summarised as the analysis of expression of the gene PITX2 - may serve as a method for predicting the survival of a patient with breast cancer. Especially the invention relates to methods for predicting the survival of a patient with breast cancer who is treated with at least one adjuvant endocrine treatment, wherein endocrine treatment is meant to comprise any treatment targeting the estrogen receptor pathway or estrogen synthesis pathway or estrogen conversion pathway i.e., which is involved in estrogen metabolism, production or secretion.

In this context survival is meant to describe the time from diagnosis or start of treatment to an endpoint, which may be the time of death (considering any reason for death or only death from breast cancer), or the time of recurrence of breast cancer, which may be local or distant, or the time of occurrence of any breast cancer associated disease. Therefore "predicting the survival" is meant to comprise predicting the disease free survival, as well as the overall survival or any other consideration of time between diagnosis and endpoint of treatment.

PITX2 (also known as PTX2, RS, RGS, ARP1, Brx1, IDG2, IGDS, IHG2, RIEG, IGDS2, IRID2, Otlx2, RIEG1, MGC20144) is known to belong to the PTX subfamily of PTX1, PTX2, and PTX3 genes which define a novel family of transcription factors, within the paired-like class of homeodomain factors. The gene PITX2 (according to NM_153426) encodes the paired-like homeodomain transcription factor 2, which is known to be expressed during development of anterior structures such as the eye, teeth, and anterior pituitary.

Toyota et al., (2001) (Blood 97: p 2823-9.) found hypermethylation of the PITX2 gene in a large proportion of acute myeloid leukemias. Furthermore, in this study hypermethylation of PITX2 is positively correlated to methylation of the ER gene and to a reduced expression level. Means to analyse the methylation pattern of the PITX2 gene have been described in a number of patent applications, too (WO 02/077272 is related to the use of methylation markers to differentiate between AML and ALL, WO 01/19845 is related to several differentially methylated sequences useful for diagnosis of several cell proliferative disorders, WO 02/00927 and WO 01/092565 are related to the use of methylation markers to diagnose diseases associated with development genes or associated with DNA transcription, respectively.

Although the methylation of PITX2 has been associated with development, transcription and disease such as cancer, it has no heretofore recognised role in the outcome prediction of breast cancer patients or responsiveness to endocrine treatment.

EXPRESSION ANALYSIS

The expression of a gene, or rather the protein encoded by the gene, can be studied on four different levels: firstly, protein expression levels can be determined directly, secondly, mRNA transcription levels can be determined, thirdly, epigenetic modifications, such as gene's DNA methylation profile or the gene's histone profile; can be analysed, as methylation is often correlated with inhibited protein expression, and fourth, the gene itself may be analysed for genetic modifications such as mutations, deletions, polymorphisms etc. influencing the expression of the gene product.

The levels of observation that have been studied by the methodological developments of recent years in molecular biology, are the genes themselves, the transcription of these genes into RNA, and the translation into the resulting proteins. However how the activation and inhibition of specific genes, in specific cells and tissues, at specific time points in the course of development of an individual are controlled, is correlatable to the degree and character of the methylation of the genes or respectively the genome. In this respect, pathogenic conditions may manifest themselves in a changed methylation pattern of individual genes or of the genome.

The four terms that apply to the fields of overall genome-wide analysis of all these biological processes are called: Proteomics, Transcriptomics, Epigenomics (or Methylomics) and Genomics. Methods and techniques that can be used for studying expression or studying the modifications responsible for expression on all of these levels are well described in the literature and therefore known to a person skilled in the art. They are described in text books of molecular biology and in a large number of scientific journals.

How to analyse the protein expression of a single gene is prior art. It usually requires an antibody specific for the gene product of interest. Appropriate technologies would be ELISA or Immunohistochemistry.

The analysis of the level of mRNA also has been described sufficiently. These days the gold standard is the reverse transcriptase PCR.

To avoid duplication a more detailed description of the prior art relating to existing and well known technologies is given within the description of the invention, as it is part of the invention.

US patent application 2003/0198970 by Gareth Roberts lists some of the technologies and methods on how to determine a person's "genetic make up", i.e. the genetic modifications, such as deletions, polymorphisms, mutations etc. that may vary between individuals and describes the potential role of this genetic sequence information in the individual's variability in disease, response to therapy and prognosis. Epigenetic differences however are not mentioned. The gene PITX2 is listed within this application as one gene name out of a long and comprehensive list of about 2.500 other gene names, suggesting its expression could play a role in some kind of treatment response. However, this is simply an assumption based on speculation only, as no experiments are disclosed, which demonstrate any kind of relation between genetic modifications of PITX2 and an individual's variation in treatment response.

A less established area in this context is the field of epigenomics or epigenetics, i.e. the field concerned with analysis of DNA methylation patterns.

Methylation of DNA can play an important role in the control of gene expression in mammalian cells. DNA methyltransferases are involved in DNA methylation and catalyse the

transfer of a methyl group from S-adenosylmethionine to cytosine residues to form 5-methylcytosine, a modified base that is found mostly at CpG sites in the genome. The presence of methylated CpG islands in the promoter region of genes can suppress their expression. This process may be due to the presence of 5-methylcytosine, which apparently interferes with the binding of transcription factors or other DNA-binding proteins to block transcription. In different types of tumours, aberrant or accidental methylation of CpG islands in the promoter region has been observed for many cancer-related genes, resulting in the silencing of their expression. Such genes include tumour suppressor genes, genes that suppress metastasis and angiogenesis, and genes that repair DNA (Mompalmer and Bovenzi (2000) J. Cell Physiol. 183:145-54).

In addition it has been described that DNA methylation may also play a role in the field of pharmacogenetics. A similar approach on how to apply information about genetic modifications of the genome to the analysis of individual responses to treatment as was for example described by Gareth Roberts in US application 2003/0198970 was already subject of the application WO 02/037398, tailored to the application of information about *epigenetic* modifications of the genome, based on DNA methylation analysis, to guide treatment selection and to study individual's treatment responses.

An example for the applicability of this idea was given by Esteller et al. (Esteller et al. (2000) N Engl J Med. 2000 Nov 9;343(19):1350-4.), who demonstrated that methylation of the MGMT promoter in gliomas is a useful predictor of the responsiveness of the tumours to alkylating agents. More recently, Frühwald has summarised a series of studies demonstrating that DNA methylation is associated with the aggressiveness of different cancers (Frühwald MC. DNA methylation patterns in cancer: novel prognostic indicators? Am J Pharmacogenomics. 2003;3(4):245-60).

An example for the potential of analysis of epigenetic modifications, such as DNA methylation analysis, for the prediction of treatment response - related to breast cancer- was presented as a poster by Martens et al. at the San Antonio Breast Cancer Symposium, San Antonio, TX, December, 3-6, 2003. Breast cancer patients which have had their tumours removed by surgery and developed metastases at some point after the removal, were treated with Tamoxifen, an endocrine treatment drug. The primary tumour samples were analysed for aberrant methylation patterns. The patients were then divided into two sub classes according to their objec-

tive tumour response: patients with progressive disease (which could be described as increasing metastasis size) and patients with complete or partial remission of the relapsed tumour (which could be described as decreasing metastasis size). It turned out, that those patients which had a tumour removed and experienced a remission (decrease in size) of the metastasis under endocrine treatment, had suffered from a tumour which showed a distinct pattern of DNA methylation at specific CpG sites, whereas patients which show progressive disease (did not experience a decrease but an increase in size of their metastases), under endocrine treatment, suffered from a tumour which did not show this distinct pattern of DNA methylation (but a different pattern) at these CpG sites. This is a clear indication, that the methylation pattern described in that study can serve as a **predictive** treatment response tool for an endocrine treatment, like tamoxifen. The results of this study, i.e. predictive biomarkers and assays therefore, are subject of patent application WO 04/035803, published at April 29, 2004: Method and nucleic acid for the improved treatment of breast cell proliferative disorders. Predictive markers as described above will also be called 'metastatic' markers in the context of this application. PITX2 is also listed as a predictive marker in said application.

Currently several predictive markers are under evaluation. As up to now most patients have received Tamoxifen as endocrine treatment most of the markers have been shown to be associated with response or resistance to tamoxifen. However, it is generally assumed that there is a large overlap between responders to one or the other endocrine treatment. In fact, ER and PR expression are used to select patients for any endocrine treatment. Among the markers which have been associated with tamoxifen response is bcl-2. High bcl-2 expression levels showed promising correlation to tamoxifen therapy response in patients with metastatic disease and prolonged survival and added valuable information to an ER negative patient subgroup (J Clin Oncology, 1997, 15 5: 1916-1922; Endocrine, 2000, 13(1):1-10). There is conflicting evidence regarding the independent predictive value of c-erbB2 (Her2/neu) overexpression in patients with advanced breast cancer that require further evaluation and verification (British J of Cancer, 1999, 79 (7/8):1220-1226; J Natl Cancer Inst, 1998, 90 (21): 1601-1608).

Other predictive markers include SRC-1 (steroid receptor coactivator-1), CGA mRNA over expression, cell kinetics and S phase fraction assays (Breast Cancer Res and Treat, 1998, 48:87-92; Oncogene, 2001, 20:6955-6959). Recently, uPA (Urokinase-type plasminogen activator) and PAI-1 (Plasminogen activator inhibitor type 1) together showed to be useful to

define a subgroup of patients who have worse prognosis and who would benefit from adjuvant systemic therapy (J Clinical Oncology, 2002, 20 n° 4). However, all of these markers need further evaluations in prospective trials as none of them is yet a validated marker of response.

Also recently published was a study related to the **prognostic** power of methylation analysis in breast cancer patients. Müller et al. (Muller HM, Widschwendter A, Fiegl H, Ivarsson L, Goebel G, Perkmann E, Marth C, Widschwendter M. (2003) DNA methylation in serum of breast cancer patients: an independent prognostic marker. Cancer Res. 2003 Nov 15; 63(22): 7641-5.) reported about a set of genes, which can be used as biomarkers in patient pre-therapeutic sera for the prognosis of breast cancer. Specific aberrant methylation patterns of two genes found in DNA from pretreatment serum of cancer patients indicated whether their prognosis was good or bad. The DNA analysed was not tumour DNA but serum DNA. Most likely the presence of a tumour-specific pattern indicates that tumour derived DNA is present, however, the absence of a specific methylation pattern, may be due to a tumour which does not show this methylation pattern, or a tumour which does not shed sufficient DNA into the blood stream. Good or bad prognosis was defined as long or short "overall survival" after surgery, without adjuvant treatment. This result therefore relates to untreated patients, only.

These 'prognostic' markers are able to answer the question whether or not a breast cancer patient should get an aggressive adjuvant treatment like chemotherapy after removal of the tumour to avoid recurrence of cancer, i.e. occurrence of metastases.

However, none of these study results and none of these markers is able to answer the specific question raised above, whether or not a breast cancer patient should get adjuvant chemotherapy after removal of the tumour to avoid recurrence of cancer, i.e. occurrence of metastases in addition or alternative to endocrine treatment (with a drug like tamoxifen, or aromatase inhibitors).

A marker for a bad prognosis for cancer patients (without treatment), might not be applicable to a patient under adjuvant treatment with a drug like tamoxifen. Therefore the test would not be able to help deciding, whether chemotherapy, including all its side effects and inherent risks, is necessary or whether endocrine treatment is sufficient, because an endocrine treatment might change the prognosis from "bad" to "good".

The predictive 'metastatic' marker set described above, would be able to identify amongst all patients which relapsed (developed metastases after surgery) those patients, which do not respond to endocrine treatment (by partial or complete remission of relapsed tumour). These markers however, cannot be applied to answer the question whether metastases will occur at all (after surgery of the primary tumour under endocrine treatment), and consequently whether it is advised to give adjuvant chemotherapy to avoid recurrence of cancer (i.e. relapse or occurrence of metastases).

In one aspect the present invention provides a marker, PITX2 (which shall be recognised as the gene encoding for the protein PITX2; according to NM_153426), that can be used to answer that question and help guiding the decision whether or not an adjuvant chemotoxic therapy shall be subscribed in addition or instead of treatment with endocrines, like tamoxifen. A marker able to answer this question will also be called 'adjuvant' marker, in the context of this application.

In addition study results presented by Paik et al. at the San Antonio Breast Cancer Symposium, San Antonio, TX, December, 3-6, 2003 provide an answer to this question, by analysing the mRNA expression pattern of 16 genes plus 5 controls with RT-PCR.

The provided invention however has the advantage that looking at only one gene or a small selection of three to five genes will give sufficient information for a validated prognosis.

For demonstration : The 'metastatic' test (use of a 'metastatic' marker) tells a patient whether she is unlikely to respond to endocrine treatment when she develops metastases. But she does not know how high the likelihood is, that she will experience a relapse at all. The 'prognostic' test (use of a 'prognostic' marker) tells a patient whether she will have a good or bad prognosis without any treatment. Even with a "bad prognosis" endocrine treatment might be enough though. The prognostic markers are not necessarily able to predict the outcome under endocrine treatment. The 'adjuvant test' (use of an 'adjuvant' marker) tells her whether she will or will not develop recurrence, without chemotherapy, even when treated with the standard -low side effected- endocrine treatment.

This invention relates to the use of PITX2, as an 'adjuvant marker', which also serves as a 'prognostic marker', especially in hormone receptor negative women, which would not get any endocrine treatment at all.

5-methylcytosine is the most frequent covalent base modification in the DNA of eukaryotic cells. It plays a role, for example, in the regulation of the transcription, in genetic imprinting, and in tumorigenesis. Therefore, the identification of 5-methylcytosine as a component of genetic information is of considerable interest. However, 5-methylcytosine positions cannot be identified by sequencing since 5-methylcytosine has the same base pairing behaviour as cytosine. Moreover, the epigenetic information carried by 5-methylcytosine is completely lost during PCR amplification.

A relatively new and currently the most frequently used method for analysing DNA for 5-methylcytosine is based upon the specific reaction of bisulfite with cytosine which, upon subsequent alkaline hydrolysis, is converted to uracil which corresponds to thymidine in its base pairing behaviour. However, 5-methylcytosine remains unmodified under these conditions. Consequently, the original DNA is converted in such a manner that methylcytosine, which originally could not be distinguished from cytosine by its hybridisation behaviour, can now be detected as the only remaining cytosine using "normal" molecular biological techniques, for example, by amplification and hybridisation or sequencing. All of these techniques are based on base pairing which can now be fully exploited. In terms of sensitivity, the prior art is defined by a method which encloses the DNA to be analysed in an agarose matrix, thus preventing the diffusion and renaturation of the DNA (bisulfite only reacts with single-stranded DNA), and which replaces all precipitation and purification steps with fast dialysis (Olek A, Oswald J, Walter J. A modified and improved method for bisulphite based cytosine methylation analysis. *Nucleic Acids Res.* 1996 Dec 15;24(24):5064-6). Using this method, it is possible to analyse individual cells, which illustrates the potential of the method. However, currently only individual regions of a length of up to approximately 3000 base pairs are analysed, a global analysis of cells for thousands of possible methylation events is not possible. However, this method cannot reliably analyse very small fragments from small sample quantities either. These are lost through the matrix in spite of the diffusion protection.

An overview of the further known methods of detecting 5-methylcytosine may be gathered from the following review article: Rein, T., DePamphilis, M. L., Zorbas, H., *Nucleic Acids Res.* 1998, 26, 2255.

To date, barring few exceptions (e.g., Zeschnigk M, Lich C, Buiting K, Doerfler W, Horsthemke B. A single-tube PCR test for the diagnosis of Angelman and Prader-Willi syndrome based on allelic methylation differences at the SNRPN locus. *Eur J Hum Genet.* 1997 Mar-Apr;5(2):94-8) the bisulfite technique is only used in research. Always, however, short, specific fragments of a known gene are amplified subsequent to a bisulfite treatment and either completely sequenced (Olek A, Walter J. The pre-implantation ontogeny of the H19 methylation imprint. *Nat Genet.* 1997 Nov;17(3):275-6) or individual cytosine positions are detected by a primer extension reaction (Gonzalzo ML, Jones PA. Rapid quantitation of methylation differences at specific sites using methylation-sensitive single nucleotide primer extension (Ms-SNuPE). *Nucleic Acids Res.* 1997 Jun 15;25(12):2529-31, WO 95/00669) or by enzymatic digestion (Xiong Z, Laird PW. COBRA: a sensitive and quantitative DNA methylation assay. *Nucleic Acids Res.* 1997 Jun 15;25(12):2532-4). In addition, detection by hybridisation has also been described (Olek et al., WO 99/28498).

Further publications dealing with the use of the bisulfite technique for methylation detection in individual genes are: Grigg G, Clark S. Sequencing 5-methylcytosine residues in genomic DNA. *Bioessays.* 1994 Jun;16(6):431-6, 431; Zeschnigk M, Schmitz B, Dittrich B, Buiting K, Horsthemke B, Doerfler W. Imprinted segments in the human genome: different DNA methylation patterns in the Prader-Willi/Angelman syndrome region as determined by the genomic sequencing method. *Hum Mol Genet.* 1997 Mar;6(3):387-95; Feil R, Charlton J, Bird AP, Walter J, Reik W. Methylation analysis on individual chromosomes: improved protocol for bisulphite genomic sequencing. *Nucleic Acids Res.* 1994 Feb 25;22(4):695-6; Martin V, Ribieras S, Song-Wang X, Rio MC, Dante R. Genomic sequencing indicates a correlation between DNA hypomethylation in the 5' region of the pS2 gene and its expression in human breast cancer cell lines. *Gene.* 1995 May 19;157(1-2):261-4; WO 97/46705 and WO 95/15373.

An overview of the Prior Art in oligomer array manufacturing can be gathered from a special edition of *Nature Genetics* (*Nature Genetics Supplement*, Volume 21, January 1999), published in January 1999; and from the literature cited therein.

Fluorescently labelled probes are often used for the scanning of immobilised DNA arrays. The simple attachment of Cy3 and Cy5 dyes to the 5'-OH of the specific probe are particularly suitable for fluorescence labels. The detection of the fluorescence of the hybridised probes may be carried out, for example via a confocal microscope. Cy3 and Cy5 dyes, besides many others, are commercially available.

Matrix Assisted Laser Desorption Ionization Mass Spectrometry (MALDI-TOF) is a very efficient development for the analysis of biomolecules (Karas M, Hillenkamp F. Laser desorption ionization of proteins with molecular masses exceeding 10,000 daltons. *Anal Chem.* 1988 Oct 15;60(20):2299-301). An analyte is embedded in a light-absorbing matrix. The matrix is evaporated by a short laser pulse thus transporting the analyte molecule into the vapour phase in an unfragmented manner. The analyte is ionised by collisions with matrix molecules. An applied voltage accelerates the ions into a field-free flight tube. Due to their different masses, the ions are accelerated at different rates. Smaller ions reach the detector sooner than bigger ones.

MALDI-TOF spectrometry is excellently suited to the analysis of peptides and proteins. The analysis of nucleic acids is somewhat more difficult (Gut I G, Beck S. DNA and Matrix Assisted Laser Desorption Ionization Mass Spectrometry. *Current Innovations and Future Trends.* 1995, 1; 147-57). The sensitivity to nucleic acids is approximately 100 times worse than to peptides and decreases disproportionally with increasing fragment size. For nucleic acids having a multiply negatively charged backbone, the ionisation process via the matrix is considerably less efficient. In MALDI-TOF spectrometry, the selection of the matrix plays an eminently important role. For the desorption of peptides, several very efficient matrixes have been found which produce a very fine crystallisation. There are now several responsive matrixes for DNA, however, the difference in sensitivity has not been reduced. The difference in sensitivity can be reduced by chemically modifying the DNA in such a manner that it becomes more similar to a peptide. Phosphorothioate nucleic acids in which the usual phosphates of the backbone are substituted with thiophosphates can be converted into a charge-neutral DNA using simple alkylation chemistry (Gut IG, Beck S. A procedure for selective DNA alkylation and detection by mass spectrometry. *Nucleic Acids Res.* 1995 Apr 25;23(8):1367-73). The coupling of a charge tag to this modified DNA results in an increase in sensitivity to the same level as that found for peptides. A further advantage of charge tag-

ging is the increased stability of the analysis against impurities which make the detection of unmodified substrates considerably more difficult.

Genomic DNA is obtained from DNA of cell, tissue or other test samples using standard methods. This standard methodology is found in references such as Sambrook, Fritsch and Maniatis eds., *Molecular Cloning: A Laboratory Manual*, 1989.

DESCRIPTION

Characterisation of a breast cancer in terms of its predicted aggressiveness enables the physician to make an informed decision as to a therapeutic regimen with appropriate risk and benefit trade offs to the patient. Aggressiveness is taken to mean one or more of decreased patient survival or disease- or relapse-free survival, increased tumour-related complications and faster progression of tumour or metastases. According to the aggressiveness of the disease an appropriate treatment or treatments may be selected from the group consisting of chemotherapy, radiotherapy, surgery, biological therapy, immunotherapy, antibody treatments, treatments involving molecularly targeted drugs, estrogen receptor modulator treatments, estrogen receptor down-regulator treatments, aromatase inhibitors treatments, ovarian ablation, treatments providing LHRH analogues or other centrally acting drugs influencing estrogen production. Wherein a cancer is characterised as 'aggressive' it is particularly preferred that a treatment such as, but not limited to, chemotherapy is provided in addition to or instead of an endocrine targeting therapy.

Using the methods and nucleic acids described herein, statistically significant models of patient disease free survival or metastasis free survival or overall survival and/or disease progression can be developed and utilised to assist patients and clinicians in determining suitable treatment options to be included in the therapeutic regimen.

In one aspect the described method is to be used to assess the utility of therapeutic regimens comprising one or more treatments which is either an aggressive therapy such as chemotherapy or a treatment which targets the estrogen receptor pathway or is involved in estrogen metabolism, production or secretion as a therapy for patients suffering from a cell proliferative disorder of the breast tissues. In particular this aspect of the method enables the physician to determine which treatments may be used in addition to or instead of said endocrine treatment.

In a further aspect the described method enables the characterisation of the cell proliferative disorder in terms of aggressiveness, thereby enabling the physician to recommend suitable treatments. Thus, the present invention will be seen to reduce the problems associated with present breast cell proliferative disorder treatment response prediction methods.

Using the methods and nucleic acids as described herein, patient survival can be evaluated before or during treatment for a cell proliferative disorder of the breast tissues, in order to provide critical information to the patient and clinician as to the likely progression of the disease. It will be appreciated, therefore, that the methods and nucleic acids exemplified herein can serve to improve a patient's quality of life and odds of treatment success by allowing both patient and clinician a more accurate assessment of the patient's treatment options.

The method according to the definition may be used for the improved treatment of all breast cell proliferative disorder patients, both pre and post menopausal and independent of their node or estrogen receptor status. However, it is particularly preferred that said patients are node-negative and estrogen receptor positive.

The present invention makes available a method for the improved treatment and monitoring of breast cell proliferative disorders, by enabling the accurate prediction of a patient's survival without systemic therapy or with endocrine therapy comprising one or more treatments which target the estrogen receptor pathway or are involved in estrogen metabolism, production, or secretion.

In a particularly preferred embodiment, the method according to the invention enables the differentiation between patients who have a high risk of relapse under said endocrine therapy and those who have a low risk of relapse under said therapy. The method particularly preferably enables the determination of a methylation pattern characteristic for a predicted survival time, in addition to the characterisation of tumours in terms of aggressiveness.

The method according to the invention may be used for the analysis of a wide variety of cell proliferative disorders of the breast tissues including, but not limited to, ductal carcinoma *in situ*, invasive ductal carcinoma, invasive lobular carcinoma, lobular carcinoma *in situ*, comedocarcinoma, inflammatory carcinoma, mucinous carcinoma, scirrhous carcinoma, colloid carcinoma, tubular carcinoma, medullary carcinoma, metaplastic carcinoma, and papillary

carcinoma and papillary carcinoma *in situ*, undifferentiated or anaplastic carcinoma and Paget's disease of the breast.

The method according to the invention is particularly suited to the prediction of survival of breast cancer in the following treatment setting. In one embodiment, the method is applied to patients who receive endocrine pathway targeting treatment as secondary treatment to an initial non chemotherapeutical therapy, e.g. surgery (hereinafter referred to as the adjuvant setting) as illustrated in Figure 1. Such a treatment is often prescribed to patients suffering from Stage 1 to 3 breast carcinomas. In this embodiment patients survival times are predicted according to their gene expression or genetic or epigenetic modifications. By detecting patients with worse disease free survival times the physician may choose to recommend the patient for further treatment, instead of or in addition to the endocrine targeting therapy(s), in particular but not limited to, chemotherapy.

This invention specifically relates to a new biomarker, PITX2, for patients diagnosed with breast cell proliferative disease, allowing the prediction of survival (or outcome) without treatment, or with different therapies, like a cytotoxic therapy (chemotherapy) in addition or instead of (for example in hormone receptor negative patients) an endocrine treatment, like treatment with Tamoxifen or aromatase inhibitors, wherein the prediction is based on the patient's survival or clinical or pathological tumour response, or response measured with other surrogate parameters.

It is also an embodiment of the invention to base the prediction of survival on a combination of different markers together with PITX2. To increase the likelihood of a correct prognosis it is preferred that the expression of some selected additional genes is investigated in addition to the analysis of PITX2 expression. It is preferred that a mini panel comprising of one or more, up to 7 additional genes is used. The preferred genes to be added to such a mini panel, would be one or more out of the group comprising ABCA8, CDK6, ERBB2, ONECUT2, PLAU, TBC1D3 and TFF1. Especially preferred would be a combined analysis of PITX2 with TBC1D3. Another preferred combination would comprise of PITX2 analysis with TBC1D3 and CDK6 analysis. Wherever in the following the invention is described specifically for PITX2, it is meant to also include a preferred combination of PITX2 with one or more of the named genes.

This invention therefore relates to new methods or tools, for patients diagnosed with breast cell proliferative disease, allowing the evaluation of adjuvant therapy based on a prediction of outcome.

More specifically this invention provides new methods or tools, for patients diagnosed with breast cell proliferative disease, allowing the evaluation of adjuvant therapy, i.e. therapy before or after surgical removal of the tumour, like a cytotoxic therapy (chemotherapy) in addition to or instead of (for example in hormone receptor negative patients) an endocrine treatment, like treatment with Tamoxifen or aromatase inhibitors, wherein the evaluation is based on the prediction of the patient's survival.

One aspect of the invention is the provision of tools for predicting the survival of a patient diagnosed with a breast cell proliferative disease, such as breast cancer. These tools comprise methods for the analysis of either the expression levels of PITX2 protein, or PITX2 mRNA or the analysis of the patient's individual genetic or epigenetic modification of the gene PITX2 – summarised as the analysis of expression of the gene PITX2. Preferably the invention relates to methods for predicting the survival of a patient diagnosed with breast cancer. Preferably said patient is treated with at least one adjuvant endocrine treatment, wherein endocrine treatment is meant to comprise any treatment targeting the estrogen receptor pathway or estrogen synthesis pathway or estrogen conversion pathway i.e., which is involved in estrogen metabolism, production or secretion. Preferably the patient was treated with said adjuvant endocrine treatment after surgical removal of the tumour. Also preferably the survival is the disease free survival.

Especially preferred are methods applied for the prediction of the disease free survival of a patient diagnosed with breast cancer under adjuvant endocrine treatment after surgical tumour removal. Even more preferred are those methods, which analyse the DNA methylation profile of the genomic region associated with the gene PITX2. Especially preferred is the analysis of the DNA methylation profile of the genomic sequence given in SEQ ID 1. Especially preferred is furthermore the analysis of the methylation status of eight specific CpG dinucleotides, covered in the three subsequences of said SEQ ID 1 given in SEQ ID NOS: 13, 18 and 19. The use of nucleic acids hybridising to these nucleic acid sequences for the prediction of survival according to the invention are preferred embodiments of said invention. The use of nucleic acids hybridising to CpG positions within these nucleic acid sequences after these

nucleic acids have been contacted with one or more agents that convert cytosine bases that are unmethylated at the 5'-position thereof to a base that is detectably dissimilar to cytosine in terms of hybridisation properties, for the prediction of survival according to the invention are especially preferred embodiments of said invention.

This methodology presents further improvements over the state of the art in that the method may be applied to any subject, independent of the estrogen and/or progesterone receptor status. Therefore in a preferred embodiment, the subject is not required to have been tested for estrogen or progesterone receptor status.

The object of the invention is preferably achieved by means of the analysis of the methylation pattern of PITX2 and/or its regulatory region. In a particularly preferred embodiment the sequence of said gene comprises SEQ ID 1 and the sequence complementary thereto.

In one preferred embodiment the object of the invention is the prediction of survival of a subject under a treatment which targets the estrogen receptor pathway or is involved in estrogen metabolism, production or secretion. This is achieved by analysis of the expression pattern of PITX2 and wherein it is further preferred that the sequence of said gene comprises SEQ ID NO: 1 or parts thereof.

In one aspect the invention discloses novel methods utilising the gene PITX2 for the prediction of survival of a patient diagnosed with a breast cell proliferative disease. In a preferred embodiment said patient diagnosed with a breast cell proliferative disease is treated with adjuvant endocrine monotherapy.

The invention discloses the use of the gene PITX2, as well as its promoter and regulatory elements as a prognostic marker for survival of breast cancer patients. It is preferred that these patients are treated with adjuvant endocrine monotherapy. More specifically, the disclosed matter shows the applicability of said gene to answer the question and help guiding the decision whether or not an adjuvant chemotoxic therapy shall be prescribed, preferably in addition to endocrine treatment, like the treatment with tamoxifen or aromatase inhibitors.

In one aspect of the invention, the disclosed matter provides novel nucleic acid sequences useful for the analysis of methylation within said gene, other aspects provide novel uses of the

gene and the gene product as well as methods, assays and kits directed to prognosing the survival of a patient diagnosed with breast cell proliferative disease. Preferably said patient is treated with adjuvant endocrine monotherapy.

In one embodiment the method discloses the use of the gene PITX2 as a marker for the prognosis of the survival of a patient suffering from a breast cell proliferative disease. Preferably said patient is treated with adjuvant endocrine monotherapy. Said use of the gene may be enabled by means of any analysis of the expression of the gene, by means of mRNA expression analysis or protein expression analysis or by analysis of its genetic modifications leading to an altered expression. However, in the most preferred embodiment of the invention, prediction of the survival of a patient diagnosed with breast cell proliferative disease, preferably treated with adjuvant endocrine monotherapy, is enabled by means of analysis of the methylation status of CpG sites within the gene PITX2 and its promoter or regulatory elements.

To detect the levels of mRNA encoding PITX2 in a detection system for breast cancer relapse, a sample is obtained from a patient. Said obtaining of a sample is not meant to be retrieving of a sample, as in performing a biopsy, but rather directed to the availability of an isolated biological material representing a specific tissue, relevant for the intended use. The sample can be a tumour tissue sample from the surgically removed tumour, a biopsy sample as taken by a surgeon and provided to the analyst or a sample of blood, plasma, serum or the like. The sample may be treated to extract the nucleic acids contained therein. The resulting nucleic acid from the sample is subjected to gel electrophoresis or other separation techniques. Detection involves contacting the nucleic acids and in particular the mRNA of the sample with a DNA sequence serving as a probe to form hybrid duplexes. The stringency of hybridisation is determined by a number of factors during hybridisation and during the washing procedure, including temperature, ionic strength, length of time and concentration of formamide. These factors are outlined in, for example, Sambrook et al. (Molecular Cloning: A Laboratory Manual, 2nd ed., 1989). Detection of the resulting duplex is usually accomplished by the use of labelled probes. Alternatively, the probe may be unlabeled, but may be detectable by specific binding with a ligand which is labelled, either directly or indirectly. Suitable labels and methods for labelling probes and ligands are known in the art, and include, for example, radioactive labels which may be incorporated by known methods (e.g., nick translation or kinasing), biotin, fluorescent groups, chemiluminescent groups (e.g., dioxetanes, particularly triggered dioxetanes), enzymes, antibodies, and the like.

In order to increase the sensitivity of the detection in a sample of mRNA encoding PITX2, the technique of reverse transcription/polymerisation chain reaction can be used to amplify cDNA transcribed from mRNA encoding PITX2. The method of reverse transcription/PCR is well known in the art (for example, see Watson and Fleming, supra).

The reverse transcription/PCR method can be performed as follows. Total cellular RNA is isolated by, for example, the standard guanidium isothiocyanate method and the total RNA is reverse transcribed. The reverse transcription method involves synthesis of DNA on a template of RNA using a reverse transcriptase enzyme and a 3' end primer. Typically, the primer contains an oligo(dT) sequence. The cDNA thus produced is then amplified using the PCR method and PITX2 specific primers. (Belyavsky et al, Nucl Acid Res 17:2919-2932, 1989; Krug and Berger, Methods in Enzymology, Academic Press, N.Y., Vol.152, pp. 316-325, 1987 which are incorporated by reference)

The present invention may also be described in certain embodiments as a kit for use in predicting the survival of a breast cancer patient before or after surgical tumour removal with or without adjuvant endocrine monotherapy state through testing of a biological sample. A representative kit may comprise one or more nucleic acid segments as described above that selectively hybridise to PITX2 mRNA and a container for each of the one or more nucleic acid segments. In certain embodiments the nucleic acid segments may be combined in a single tube. In further embodiments, the nucleic acid segments may also include a pair of primers for amplifying the target mRNA. Such kits may also include any buffers, solutions, solvents, enzymes, nucleotides, or other components for hybridisation, amplification or detection reactions. Preferred kit components include reagents for reverse transcription-PCR, in situ hybridisation, Northern analysis and/or RPA.

The present invention further provides for methods to detect the presence of the polypeptide, PITX2, in a sample obtained from a patient. It is preferred that said sequence is essentially the same as the sequence presented in SEQ ID 20, as given in figure 10. Any method known in the art for detecting proteins can be used. Such methods include, but are not limited to immunodiffusion, immunoelectrophoresis, immunochemical methods, binder-ligand assays, immunohistochemical techniques, agglutination and complement assays. (for example see Basic and Clinical Immunology, Sites and Terr, eds., Appleton & Lange, Norwalk, Conn. pp 217-

262, 1991 which is incorporated by reference). Preferred are binder-ligand immunoassay methods including reacting antibodies with an epitope or epitopes of PITX2 and competitively displacing a labelled PITX2 protein or derivative thereof.

Certain embodiments of the present invention comprise the use of antibodies specific to the polypeptide encoded by the PITX2 gene. Such antibodies may be useful for prognosing the survival of a breast cancer patient preferably under adjuvant endocrine monotherapy by comparing a patient's levels of PITX2 marker expression to expression of the same marker in normal individuals. In certain embodiments production of monoclonal or polyclonal antibodies can be induced by the use of the PITX2 polypeptide as antigen. Such antibodies may in turn be used to detect expressed proteins as markers for prognosis of relapse of a breast cancer patient under adjuvant endocrine monotherapy. The levels of such proteins present in the peripheral blood of a patient may be quantified by conventional methods. Antibody-protein binding may be detected and quantified by a variety of means known in the art, such as labeling with fluorescent or radioactive ligands. The invention further comprises kits for performing the above-mentioned procedures, wherein such kits contain antibodies specific for the PITX2 polypeptides.

Numerous competitive and non-competitive protein binding immunoassays are well known in the art. Antibodies employed in such assays may be unlabeled, for example as used in agglutination tests, or labelled for use a wide variety of assay methods. Labels that can be used include radionuclides, enzymes, fluorescers, chemilumescers, enzyme substrates or co-factors, enzyme inhibitors, particles, dyes and the like for use in radioimmunoassay (RIA), enzyme immunoassays, e.g., enzyme-linked immunosorbent assay (ELISA), fluorescent immunoassays and the like. Polyclonal or monoclonal antibodies to PITX2 or an epitope thereof can be made for use in immunoassays by any of a number of methods known in the art. One approach for preparing antibodies to a protein is the selection and preparation of an amino acid sequence of all or part of the protein, chemically synthesising the sequence and injecting it into an appropriate animal, usually a rabbit or a mouse (Milstein and Kohler *Nature* 256:495-497, 1975; Gutfre and Milstein, *Methods in Enzymology: Immunochemical Techniques* 73:1-46, Langone and Banatis eds., Academic Press, 1981 which are incorporated by reference). Methods for preparation of PITX2 or an epitope thereof include, but are not limited to chemical synthesis, recombinant DNA techniques or isolation from biological samples.

The invention provides significant improvements over the state of the art in that -at the time of filing- there are no single markers known to the public which can be used to predict the likelihood of relapse or of survival of a breast cancer patient under adjuvant endocrine monotherapy, neither from tissue samples nor from body fluid samples.

Also, no methylation marker is known which can be used to detect the likelihood of relapse or of survival of a breast cancer patient. Especially, no methylation marker is known which can be used to detect the likelihood of relapse or of survival of a breast cancer patient under adjuvant endocrine monotherapy, neither from tissue samples nor from body fluid samples.

The objective of the invention is also preferably achieved by analysis of the methylation state of the CpG dinucleotides within the genomic sequence according to SEQ ID NO: 1 and sequences complementary thereto. SEQ ID NO: 1 discloses the gene PITX2 and its promoter and regulatory elements, wherein said fragment comprises CpG dinucleotides exhibiting a disease specific methylation pattern. The methylation pattern of the gene PITX2 and its promoter and regulatory elements have heretofore not been analysed with regard to prognosis or prediction of survival of a patient diagnosed with a breast cell proliferative disorder. Due to the degeneracy of the genetic code, the sequence as identified in SEQ ID NO: 1 should be interpreted so as to include all substantially similar and equivalent sequences upstream of the promoter region of a gene which encodes a polypeptide with the biological activity of that encoded by PITX2.

In a preferred embodiment of the method, the objective of the invention is achieved by analysis of a nucleic acid comprising a sequence of at least 18 bases in length according to one of SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto.

The sequences of SEQ ID NOS: 2 to 5 provide modified versions of the nucleic acid according to SEQ ID NO: 1, wherein the conversion of said sequence results in the synthesis of a nucleic acid having a sequence that is unique and distinct from SEQ ID NO: 1 as follows. (*see also* the following TABLE 1): SEQ ID NO: 1, sense DNA strand of PITX2 gene and its promoter and regulatory elements; SEQ ID NO: 2, converted SEQ ID NO: 1, wherein "C" converted to "T," but "CpG" remains "CpG." (*i.e.*, corresponds to case where, for SEQ ID NO: 1, all "C" residues of CpG dinucleotide sequences are methylated and are thus not converted); SEQ ID NO: 3, complement of SEQ ID NO: 1, wherein "C" converted to "T," but "CpG"

remains "CpG" (*i.e.*, corresponds to case where, for the complement (*antisense* strand) of SEQ ID NO: 1, all "C" residues of CpG dinucleotide sequences are methylated and are thus not converted); SEQ ID NO: 4, converted SEQ ID NO: 1, wherein "C" converted to "T" for all "C" residues, including those of "CpG" dinucleotide sequences (*i.e.*, corresponds to case where, for SEQ ID NO: 1, all "C" residues of CG dinucleotide sequences are unmethylated); SEQ ID NO: 5, complement of SEQ ID NO: 1, wherein "C" converted to "T" for all "C" residues, including those of "CpG" dinucleotide sequences (*i.e.*, corresponds to case where, for the complement (*antisense* strand) of SEQ ID NO: 1, all "C" residues of CpG dinucleotide sequences are unmethylated).

TABLE 1. Description of SEQ ID NOS: 1 to 5

SEQ ID NO	Relationship to SEQ ID NO:1	Nature of cytosine base conversion
SEQ ID NO:1	Sense strand (PITX2 gene including promoter and regulatory elements)	None; untreated sequence
SEQ ID NO:2	Converted sense strand	"C" to "T," but "CpG" remains "CpG" (all "C" residues of CpGs are methylated)
SEQ ID NO:3	Converted antisense strand	"C" to "T," but "CpG" remains "CpG" (all "C" residues of CpGs are methylated)
SEQ ID NO:4	Converted sense strand	"C" to "T" for all "C" residues (all "C" residues of CpGs are <u>un</u> methylated)
SEQ ID NO:5	Converted antisense strand	"C" to "T" for all "C" residues (all "C" residues of CpGs are <u>un</u> methylated)

Significantly, heretofore, the nucleic acid sequences and molecules according to SEQ ID NO: 1 to SEQ ID NO: 5 were not implicated in or connected with the ascertainment of the prognosis of breast cancer relapse or the prediction of survival of breast cancer patients.

The described invention further discloses oligonucleotides or oligomers for detecting the cytosine methylation state within pretreated DNA, according to SEQ ID NO: 2 to SEQ ID NO: 5. The use of said oligonucleotides or oligomers comprising a nucleic acid sequence having a length of at least nine (9) nucleotides which hybridise, under moderately stringent or stringent conditions (as defined herein above), to a pretreated nucleic acid sequence according to SEQ

ID NO: 2 to SEQ ID NO: 5 and/or sequences complementary thereto is another embodiment of this invention.

Thus, the present invention includes the use of nucleic acid molecules (*e.g.*, oligonucleotides and peptide nucleic acid (PNA) molecules (PNA-oligomers)) that hybridise under moderately stringent and/or stringent hybridisation conditions to all or a portion of the sequences of SEQ ID NOS: 2 to 5, or to the complements thereof for prediction of survival according to the invention. The hybridising portion of the hybridising nucleic acids is typically at least 9, 15, 20, 25, 30 or 35 nucleotides in length. However, longer molecules have inventive utility, and are thus within the scope of the present invention.

Preferably, the hybridising portion of the inventive hybridising nucleic acids is at least 95%, or at least 98%, or 100% identical to the sequence, or to a portion thereof of SEQ ID NOS: 2 to 5, or to the complements thereof.

Hybridising nucleic acids of the type described herein can be used, for example, as a primer (*e.g.*, a PCR primer), or a diagnostic and/or prognostic probe or primer. Preferably, hybridisation of the oligonucleotide probe to a nucleic acid sample is performed under stringent conditions and the probe is 100% identical to the target sequence. Nucleic acid duplex or hybrid stability is expressed as the melting temperature or T_m , which is the temperature at which a probe dissociates from a target DNA. This melting temperature is used to define the required stringency conditions.

For target sequences that are related and substantially identical to the corresponding sequence of SEQ ID NO: 1 (such as PITX2 allelic variants and SNPs), rather than identical, it is useful to first establish the lowest temperature at which only homologous hybridisation occurs with a particular concentration of salt (*e.g.*, SSC or SSPE). Then, assuming that 1% mismatching results in a 1°C decrease in the T_m , the temperature of the final wash in the hybridisation reaction is reduced accordingly (for example, if sequences having > 95% identity with the probe are sought, the final wash temperature is decreased by 5°C). In practice, the change in T_m can be between 0.5°C and 1.5°C per 1% mismatch.

Examples of inventive oligonucleotides of length X (in nucleotides), as indicated by polynucleotide positions with reference to, *e.g.*, SEQ ID NO: 1, include those corresponding to sets

of consecutively overlapping oligonucleotides of length X, where the oligonucleotides within each consecutively overlapping set (corresponding to a given X value) are defined as the finite set of Z oligonucleotides from nucleotide positions:

n to (n + (X-1));

where n=1, 2, 3,...(Y-(X-1));

where Y equals the length (nucleotides or base pairs) of SEQ ID NO: 1 ;

where X equals the common length (in nucleotides) of each oligonucleotide in the set (e.g., X=20 for a set of consecutively overlapping 20-mers); and

where the number (Z) of consecutively overlapping oligomers of length X for a given SEQ ID NO of length Y is equal to Y-(X-1). For example Z=2,785-19=2,766 for either sense or antisense sets of SEQ ID NO: 1, where X=20.

Preferably, the set is limited to those oligomers that comprise at least one CpG, Cpa or tpG dinucleotide, wherein 'Cpa' is indicating that said Cpa hybridises to a position (tpG) which was a CpG prior to bisulfite conversion and is a TpG now; and wherein 'tpG' is indicating that said tpG hybridises to a position (Cpa) which is the complementary to a position (tpG) which was a CpG prior to bisulfite conversion and is a TpG now.

The present invention encompasses, for *each* of SEQ ID NOS: 2 to 5 (sense and antisense), the use of multiple consecutively overlapping sets of oligonucleotides or modified oligonucleotides of length X, where, e.g., X= 9, 10, 17, 20, 22, 23, 25, 27, 30 or 35 nucleotides.

The oligonucleotides or oligomers according to the present invention constitute effective tools useful to ascertain genetic and epigenetic parameters of the genomic sequence corresponding to SEQ ID NO: 1. Preferred sets of such oligonucleotides or modified oligonucleotides of length X are those consecutively overlapping sets of oligomers corresponding to SEQ ID NOS:1-5 (and to the complements thereof). Preferably, said oligomers comprise at least one CpG, tpG or Cpa dinucleotide.

Particularly preferred oligonucleotides or oligomers used to the present invention are those in which the cytosine of the CpG dinucleotide (or of the corresponding converted TpG or CpA dinucleotide) sequences is within the middle third of the oligonucleotide; that is, where the oligonucleotide is, for example, 13 bases in length, the CpG, TpG or CpA dinucleotide is positioned within the fifth to ninth nucleotide from the 5'-end.

The oligonucleotides used in this invention can also be modified by chemically linking the oligonucleotide to one or more moieties or conjugates to enhance the activity, stability or detection of the oligonucleotide. Such moieties or conjugates include chromophores, fluorophors, lipids such as cholesterol, cholic acid, thioether, aliphatic chains, phospholipids, polyamines, polyethylene glycol (PEG), palmityl moieties, and others as disclosed in, for example, United States Patent Numbers 5,514,758, 5,565,552, 5,567,810, 5,574,142, 5,585,481, 5,587,371, 5,597,696 and 5,958,773. The probes may also exist in the form of a PNA (peptide nucleic acid) which has particularly preferred pairing properties. Thus, the oligonucleotide may include other appended groups such as peptides, and may include hybridisation-triggered cleavage agents (Krol et al., *BioTechniques* 6:958-976, 1988) or intercalating agents (Zon, *Pharm. Res.* 5:539-549, 1988). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a chromophore, fluorophor, peptide, hybridisation-triggered cross-linking agent, transport agent, hybridisation-triggered cleavage agent, etc.

The oligonucleotide may also comprise at least one art-recognised modified sugar and/or base moiety, or may comprise a modified backbone or non-natural internucleoside linkage.

The oligomers used in the present invention are normally used in so called "sets" which contain at least one oligomer for analysis of each of the CpG dinucleotides of a genomic sequence comprising SEQ ID NO: 1 and sequences complementary thereto or to their corresponding CG, tG or Ca dinucleotide within the pretreated nucleic acids according to SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, wherein a 't' indicates a nucleotide which converted from a cytosine into a thymine and wherein 'a' indicates the complementary nucleotide to such a converted thymine. Preferred is a set which contains at least one oligomer for each of the CpG dinucleotides within the gene PITX2 and its promoter and regulatory elements in both the pretreated and genomic versions of said gene, SEQ ID NO: 2 to 5 and SEQ ID NO: 1, respectively. However, it is anticipated that for economic or other factors it may be preferable to analyse a limited selection of the CpG dinucleotides within said sequences and the contents of the set of oligonucleotides should be altered accordingly. Therefore, the present invention moreover relates to a set of at least 3 n (oligonucleotides and/or PNA-oligomers) used for detecting the cytosine methylation state in pretreated genomic DNA (SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto) and genomic DNA (SEQ ID NO: 1 and sequences complementary thereto). These probes enable

diagnosis and/or therapy of genetic and epigenetic parameters of cell proliferative disorders. The set of oligomers may also be used for detecting single nucleotide polymorphisms (SNPs) in pretreated genomic DNA (SEQ ID NO: 2 to SEQ ID NO: 5, and sequences complementary thereto) and genomic DNA (SEQ ID NO: 1, and sequences complementary thereto) .

Moreover, the present invention includes the use of a set of at least two oligonucleotides which can be used as so-called "primer oligonucleotides" for amplifying DNA sequences of one of SEQ ID NO: 1 to SEQ ID NO: 5 and sequences complementary thereto, or segments thereof.

In the case of the sets of oligonucleotides according to the present invention, it is preferred that at least one and more preferably all members of the set of oligonucleotides is bound to a solid phase.

According to the present invention, it is preferred that an arrangement of different oligonucleotides and/or PNA-oligomers (a so-called "array") made available by the present invention is present in a manner that it is likewise bound to a solid phase. This array of different oligonucleotide- and/or PNA-oligomer sequences can be characterised in that it is arranged on the solid phase in the form of a rectangular or hexagonal lattice. The solid phase surface is preferably composed of silicon, glass, polystyrene, aluminium, steel, iron, copper, nickel, silver, or gold. However, nitrocellulose as well as plastics such as nylon which can exist in the form of pellets or also as resin matrices may also be used.

Therefore, a further subject matter of the present invention is a method for manufacturing an array fixed to a carrier material for analysis in connection with cell proliferative disorders, in which method at least one oligomer according to the present invention is coupled to a solid phase. Methods for manufacturing such arrays are known, for example, from US Patent 5,744,305 by means of solid-phase chemistry and photolabile protecting groups.

A further subject matter of the present invention relates to a DNA chip for the analysis of cell proliferative disorders. DNA chips are known, for example, in US Patent 5,837,832.

The described invention further provides a composition of matter useful for prognosing the relapse of breast cancer patients or predicting the survival of breast cancer patients. Said com-

position comprising at least one nucleic acid 18 base pairs in length of a segment of the nucleic acid sequence disclosed in SEQ ID NO: 2 to 5, and one or more substances taken from the group comprising :

1-5 mM Magnesium Chloride, 100-500 μ M dNTP, 0.5-5 units/10 μ l of taq polymerase, bovine serum albumen, an oligomer in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto. It is preferred that said composition of matter comprises a buffer solution appropriate for the stabilisation of said nucleic acid in an aqueous solution and enabling polymerase based reactions within said solution.. Suitable buffers are known in the art and commercially available.

The present invention further provides a method for conducting an assay in order to ascertain genetic and/or epigenetic parameters of the gene PITX2 and its promoter and regulatory elements. Most preferably the assay according to the following method is used in order to detect methylation within the gene PITX2 wherein said methylated nucleic acids are present in a solution further comprising an excess of background DNA, wherein the background DNA is present in between 100 to 1000 times the concentration of the DNA to be detected. Said method comprising contacting a nucleic acid sample obtained from said subject with at least one reagent or a series of reagents, wherein said reagent or series of reagents, distinguishes between methylated and non-methylated CpG dinucleotides within the target nucleic acid.

Preferably, said method comprises the following steps: In the first *step*, a sample of the tissue to be analysed is obtained. The source may be any suitable source, preferably, the source of the sample is selected from the group consisting of histological slides, biopsies, paraffin-embedded tissue, bodily fluids, plasma, serum, stool, urine, blood, nipple aspirate and combinations thereof. Preferably, the source is tumour tissue, biopsies, serum, urine, blood or nipple aspirate. The most preferred source, is the tumour sample, surgically removed from the patient or a biopsy sample of said patient.

The DNA is then isolated from the sample. Extraction may be by means that are standard to one skilled in the art, including the use of detergent lysates, sonification and vortexing with glass beads. Once the nucleic acids have been extracted, the genomic double stranded DNA is

used in the analysis.

In the second step of the method, the genomic DNA sample is treated in such a manner that cytosine bases which are unmethylated at the 5'-position are converted to uracil, thymine, or another base which is dissimilar to cytosine in terms of hybridisation behaviour. This will be understood as 'pretreatment' herein.

The above described pretreatment of genomic DNA is preferably carried out with bisulfite (hydrogen sulfite, disulfite) and subsequent alkaline hydrolysis which results in a conversion of non-methylated cytosine nucleobases to uracil or to another base which is dissimilar to cytosine in terms of base pairing behaviour. Enclosing the DNA to be analysed in an agarose matrix, thereby preventing the diffusion and renaturation of the DNA (bisulfite only reacts with single-stranded DNA), and replacing all precipitation and purification steps with fast dialysis (Olek A, et al., A modified and improved method for bisulfite based cytosine methylation analysis, *Nucleic Acids Res.* 24:5064-6, 1996) is one preferred example how to perform said pretreatment. It is further preferred that the bisulfite treatment is carried out in the presence of a radical scavenger or DNA denaturing agent.

In the third step of the method, fragments of the pretreated DNA are amplified. Wherein the source of the DNA is free DNA from serum, or DNA extracted from paraffin it is particularly preferred that the size of the amplificate fragment is between 100 and 200 base pairs in length, and wherein said DNA source is extracted from cellular sources (e.g. tissues, biopsies, cell lines) it is preferred that the amplificate is between 100 and 350 base pairs in length. It is particularly preferred that said amplificates comprise at least one 20 base pair sequence comprising at least three CpG dinucleotides. Said amplification is carried out using sets of primer oligonucleotides according to the present invention, and a preferably heat-stable polymerase. The amplification of several DNA segments can be carried out simultaneously in one and the same reaction vessel, in one embodiment of the method preferably six or more fragments are amplified simultaneously. Typically, the amplification is carried out using a polymerase chain reaction (PCR). The set of primer oligonucleotides includes at least two oligonucleotides whose sequences are each reverse complementary, identical, or hybridise under stringent or highly stringent conditions to an at least 18-base-pair long segment of the base sequences of SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto.

In one especially preferred embodiment of the method the primers may be selected from the group consisting of SEQ ID NO: 6 to SEQ ID NO: 10.

In an alternate embodiment of the method, the methylation status of preselected CpG positions within the nucleic acid sequences comprising SEQ ID NO: 2 to SEQ ID NO: 5 may be detected by use of methylation-specific primer oligonucleotides. This technique (MSP) has been described in United States Patent No. 6,265,171 to Herman. The use of methylation status specific primers for the amplification of bisulfite treated DNA allows the differentiation between methylated and unmethylated nucleic acids. MSP primer pairs contain at least one primer which hybridises to a bisulfite treated CpG dinucleotide. Therefore, the sequence of said primers comprises at least one CpG, TpG or CpA dinucleotide. MSP primers specific for non-methylated DNA contain a "T" at the 3' position of the C position in the CpG. Preferably, therefore, the base sequence of said primers is required to comprise a sequence having a length of at least 18 nucleotides which hybridises to a pretreated nucleic acid sequence according to SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, wherein the base sequence of said oligomers comprises at least one CpG, tpG or Cpa dinucleotide. In this embodiment of the method according to the invention it is particularly preferred that the MSP primers comprise between 2 and 4 CpG, tpG or Cpa dinucleotides. It is further preferred that said dinucleotides are located within the 3' half of the primer e.g. wherein a primer is 18 bases in length the specified dinucleotides are located within the first 9 bases from the 3' end of the molecule. In addition to the CpG, tpG or Cpa dinucleotides it is further preferred that said primers should further comprise several bisulfite converted bases (i.e. cytosine converted to thymine, or on the hybridising strand, guanine converted to adenosine). In a further preferred embodiment said primers are designed so as to comprise no more than 2 cytosine or guanine bases.

The fragments obtained by means of the amplification can carry a directly or indirectly detectable label. Preferred are labels in the form of fluorescence labels, radionuclides, or detachable molecule fragments having a typical mass which can be detected in a mass spectrometer. Where said labels are mass labels, it is preferred that the labelled amplicates have a single positive or negative net charge, allowing for better detectability in the mass spectrometer. The detection may be carried out and visualised by means of, e.g., matrix assisted laser desorption/ionisation mass spectrometry (MALDI) or using electron spray mass spectrometry (ESI).

Matrix Assisted Laser Desorption/Ionization Mass Spectrometry (MALDI-TOF) is a very efficient development for the analysis of biomolecules (Karas & Hillenkamp, *Anal Chem.*, 60:2299-301, 1988). An analyte is embedded in a light-absorbing matrix. The matrix is evaporated by a short laser pulse thus transporting the analyte molecule into the vapour phase in an unfragmented manner. The analyte is ionised by collisions with matrix molecules. An applied voltage accelerates the ions into a field-free flight tube. Due to their different masses, the ions are accelerated at different rates. Smaller ions reach the detector sooner than bigger ones. MALDI-TOF spectrometry is well suited to the analysis of peptides and proteins. The analysis of nucleic acids is somewhat more difficult (Gut & Beck, *Current Innovations and Future Trends*, 1:147-57, 1995). The sensitivity with respect to nucleic acid analysis is approximately 100-times less than for peptides, and decreases disproportionally with increasing fragment size. Moreover, for nucleic acids having a multiply negatively charged backbone, the ionisation process via the matrix is considerably less efficient. In MALDI-TOF spectrometry, the selection of the matrix plays an eminently important role. For the desorption of peptides, several very efficient matrixes have been found which produce a very fine crystallisation. There are now several responsive matrixes for DNA, however, the difference in sensitivity between peptides and nucleic acids has not been reduced. This difference in sensitivity can be reduced, however, by chemically modifying the DNA in such a manner that it becomes more similar to a peptide. For example, phosphorothioate nucleic acids, in which the usual phosphates of the backbone are substituted with thiophosphates, can be converted into a charge-neutral DNA using simple alkylation chemistry (Gut & Beck, *Nucleic Acids Res.* 23: 1367-73, 1995). The coupling of a charge tag to this modified DNA results in an increase in MALDI-TOF sensitivity to the same level as that found for peptides. A further advantage of charge tagging is the increased stability of the analysis against impurities, which makes the detection of unmodified substrates considerably more difficult.

In a particularly preferred embodiment of the method the amplification of step three is carried out in the presence of at least one species of blocker oligonucleotides. The use of such blocker oligonucleotides has been described by Yu et al., *BioTechniques* 23:714-720, 1997. The use of blocking oligonucleotides enables the improved specificity of the amplification of a subpopulation of nucleic acids. Blocking probes hybridised to a nucleic acid suppress, or hinder the polymerase mediated amplification of said nucleic acid. In one embodiment of the method blocking oligonucleotides are designed so as to hybridise to background DNA. In a further embodiment of the method said oligonucleotides are designed so as to hinder or suppress the

amplification of unmethylated nucleic acids as opposed to methylated nucleic acids or vice versa.

Blocking probe oligonucleotides are hybridised to the bisulfite treated nucleic acid concurrently with the PCR primers. PCR amplification of the nucleic acid is terminated at the 5' position of the blocking probe, such that amplification of a nucleic acid is suppressed where the complementary sequence to the blocking probe is present. The probes may be designed to hybridise to the bisulfite treated nucleic acid in a methylation status specific manner. For example, for detection of methylated nucleic acids within a population of unmethylated nucleic acids, suppression of the amplification of nucleic acids which are unmethylated at the position in question would be carried out by the use of blocking probes comprising a 'TpG' at the position in question, as opposed to a 'CpG.' In one embodiment of the method the sequence of said blocking oligonucleotides should be identical or complementary to molecule is complementary or identical to a sequence at least 18 base pairs in length selected from the group consisting of SEQ ID NOS: 2 to 5, preferably comprising one or more CpG, TpG or CpA dinucleotides. In one embodiment of the method the sequence of said oligonucleotides is selected from the group consisting SEQ ID NO: 15 and SEQ ID NO: 16 and sequences complementary thereto.

For PCR methods using blocker oligonucleotides, efficient disruption of polymerase-mediated amplification requires that blocker oligonucleotides not be elongated by the polymerase. Preferably, this is achieved through the use of blockers that are 3'-deoxyoligonucleotides, or oligonucleotides derivatised at the 3' position with other than a "free" hydroxyl group. For example, 3'-O-acetyl oligonucleotides are representative of a preferred class of blocker molecule.

Additionally, polymerase-mediated decomposition of the blocker oligonucleotides should be precluded. Preferably, such preclusion comprises either use of a polymerase lacking 5'-3' exonuclease activity, or use of modified blocker oligonucleotides having, for example, thioate bridges at the 5'-termini thereof that render the blocker molecule nuclease-resistant. Particular applications may not require such 5' modifications of the blocker. For example, if the blocker- and primer-binding sites overlap, thereby precluding binding of the primer (*e.g.*, with excess blocker), degradation of the blocker oligonucleotide will be substantially precluded. This is because the polymerase will not extend the primer toward, and through (in the 5'-3'

direction) the blocker - a process that normally results in degradation of the hybridised blocker oligonucleotide.

A particularly preferred blocker/PCR embodiment, for purposes of the present invention and as implemented herein, comprises the use of peptide nucleic acid (PNA) oligomers as blocking oligonucleotides. Such PNA blocker oligomers are ideally suited, because they are neither decomposed nor extended by the polymerase.

In one embodiment of the method, the binding site of the blocking oligonucleotide is identical to, or overlaps with that of the primer and thereby hinders the hybridisation of the primer to its binding site. In a further preferred embodiment of the method, two or more such blocking oligonucleotides are used. In a particularly preferred embodiment, the hybridisation of one of the blocking oligonucleotides hinders the hybridisation of a forward primer, and the hybridisation of another of the probe (blocker) oligonucleotides hinders the hybridisation of a reverse primer that binds to the amplificate product of said forward primer.

In an alternative embodiment of the method, the blocking oligonucleotide hybridises to a location between the reverse and forward primer positions of the treated background DNA, thereby hindering the elongation of the primer oligonucleotides.

It is particularly preferred that the blocking oligonucleotides are present in at least 5 times the concentration of the primers.

In the fourth step of the method, the amplificates obtained during the third step of the method are analysed in order to ascertain the methylation status of the CpG dinucleotides prior to the treatment.

In embodiments where the amplificates were obtained by means of MSP amplification and/or blocking oligonucleotides, the presence or absence of an amplificate is in itself indicative of the methylation state of the CpG positions covered by the primers and or blocking oligonucleotide, according to the base sequences thereof. All possible known molecular biological methods may be used for this detection, including, but not limited to gel electrophoresis, sequencing, liquid chromatography, hybridisations, real time PCR analysis or combinations thereof. This step of the method further acts as a qualitative control of the preceding steps.

In the fourth step of the method amplicates obtained by means of both standard and methylation specific PCR are further analysed in order to determine the CpG methylation status of the genomic DNA isolated in the first step of the method. This may be carried out by means of hybridisation-based methods such as, but not limited to, array technology and probe based technologies as well as by means of techniques such as sequencing and template directed extension.

In one embodiment of the method, the amplicates synthesised in step three are subsequently hybridised to an array or a set of oligonucleotides and/or PNA probes. In this context, the hybridisation takes place in the following manner: the set of probes used during the hybridisation is preferably composed of at least 2 oligonucleotides or PNA-oligomers; in the process, the amplicates serve as probes which hybridise to oligonucleotides previously bonded to a solid phase; the non-hybridised fragments are subsequently removed; said oligonucleotides contain at least one base sequence having a length of at least 9 nucleotides which is reverse complementary or identical to a segment of the base sequences specified in the SEQ ID NO: 2 to SEQ ID NO: 5; and the segment comprises at least one CpG , TpG or CpA dinucleotide.

In a preferred embodiment, said dinucleotide is present in the central third of the oligomer. For example, wherein the oligomer comprises one CpG dinucleotide, said dinucleotide is preferably the fifth to ninth nucleotide from the 5'-end of a 13-mer. One oligonucleotide exists for the analysis of each CpG dinucleotide within the sequence according to SEQ ID NO: 1, and the equivalent positions within SEQ ID NOS: 2 to 5. Said oligonucleotides may also be present in the form of peptide nucleic acids. The non-hybridised amplicates are then removed. The hybridised amplicates are detected. In this context, it is preferred that labels attached to the amplicates are identifiable at each position of the solid phase at which an oligonucleotide sequence is located.

In yet a further embodiment of the method, the genomic methylation status of the CpG positions may be ascertained by means of oligonucleotide probes that are hybridised to the bisulfite treated DNA concurrently with the PCR amplification primers (wherein said primers may either be methylation specific or standard).

A particularly preferred embodiment of this method is the use of fluorescence-based Real

Time Quantitative PCR (Heid et al., *Genome Res.* 6:986-994, 1996; also see United States Patent No. 6,331,393). There are two preferred embodiments of utilising this method. One embodiment, known as the TaqMan™ assay employs a dual-labelled fluorescent oligonucleotide probe. The TaqMan™ PCR reaction employs the use of a non-extendible interrogating oligonucleotide, called a TaqMan™ probe, which is designed to hybridise to a CpG-rich sequence located between the forward and reverse amplification primers. The TaqMan™ probe further comprises a fluorescent “reporter moiety” and a “quencher moiety” covalently bound to linker moieties (e.g., phosphoramidites) attached to the nucleotides of the TaqMan™ oligonucleotide. Hybridised probes are displaced and broken down by the polymerase of the amplification reaction thereby leading to an increase in fluorescence. For analysis of methylation within nucleic acids subsequent to bisulfite treatment, it is required that the probe be methylation specific, as described in United States Patent No. 6,331,393, (hereby incorporated by reference in its entirety) also known as the MethyLight assay. The second preferred embodiment of this MethyLight technology is the use of dual-probe technology (Lightcycler®), each probe carrying donor or recipient fluorescent moieties, hybridisation of two probes in proximity to each other is indicated by an increase or fluorescent amplification primers. Both these techniques may be adapted in a manner suitable for use with bisulfite treated DNA, and moreover for methylation analysis within CpG dinucleotides.

Also any combination of these probes or combinations of these probes with other known probes may be used.

In a further preferred embodiment of the method, the fourth step of the method comprises the use of template-directed oligonucleotide extension, such as MS-SNuPE as described by Gonzalgo & Jones, *Nucleic Acids Res.* 25:2529-2531, 1997. In said embodiment it is preferred that the methylation specific single nucleotide extension primer (MS-SNuPE primer) is identical or complementary to a sequence at least nine but preferably no more than twenty five nucleotides in length of one or more of the sequences taken from the group of SEQ ID NO: 2 to SEQ ID NO: 5. However it is preferred to use fluorescently labelled nucleotides, instead of radiolabelled nucleotides.

In yet a further embodiment of the method, the fourth step of the method comprises sequencing and subsequent sequence analysis of the amplificate generated in the third step of the method (Sanger F., et al., *Proc Natl Acad Sci USA* 74:5463-5467, 1977).

Additional embodiments of the invention provide a method for the analysis of the methylation status of genomic DNA according to the invention (SEQ ID NO: 1) without the need for pre-treatment.

In the *first step* of such additional embodiments, the genomic DNA sample is isolated from tissue or cellular sources. Preferably, such sources include cell lines, histological slides, biopsy tissue, body fluids, or breast tumour tissue embedded in paraffin. Extraction may be by means that are standard to one skilled in the art, including but not limited to the use of detergent lysates, sonification and vortexing with glass beads. Once the nucleic acids have been extracted, the genomic double-stranded DNA is used in the analysis.

In a preferred embodiment, the DNA may be cleaved prior to the treatment, and this may be by any means standard in the state of the art, but preferably with methylation-sensitive restriction endonucleases.

In the *second step*, the DNA is then digested with one or more methylation sensitive restriction enzymes. The digestion is carried out such that hydrolysis of the DNA at the restriction site is informative of the methylation status of a specific CpG dinucleotide.

In the *third step*, which is optional but a preferred embodiment, the restriction fragments are amplified. This is preferably carried out using a polymerase chain reaction, and said amplicates may carry suitable detectable labels as discussed above, namely fluorophore labels, radionuclides and mass labels.

In the *final step* the amplicates are detected. The detection may be by any means standard in the art, for example, but not limited to, gel electrophoresis analysis, hybridisation analysis, incorporation of detectable tags within the PCR products, DNA array analysis, MALDI or ESI analysis.

The present invention enables prognosis of events which are disadvantageous to patients or individuals in which important genetic and/or epigenetic parameters within the PITX2 gene and its promoter or regulatory elements may be used as prognostic markers for breast cancer relapse or as 'adjuvant marker' for prediction of need of additional treatment besides of endo-

crine monotherapy. Said parameters obtained by means of the present invention may be compared to another set of genetic and/or epigenetic parameters, the differences serving as the basis for a prognosis of events which are disadvantageous to patients or individuals.

Specifically, the present invention provides for prognostic cancer relapse assays based on measurement of differential methylation of PITX2 CpG dinucleotide sequences. Preferred gene sequences useful to measure such differential methylation are represented herein by SEQ ID NOS: 1 to 5. Typically, such assays involve obtaining a tissue sample from a test tissue, performing an assay to measure the methylation status of at least one of the inventive PITX2-specific CpG dinucleotide sequences derived from the tissue sample, relative to a control sample, and making a diagnosis or prognosis or prediction based thereon.

In particular preferred embodiments, inventive oligomers to assess PITX2 specific CpG dinucleotide methylation status, such as those based on SEQ ID NOS: 1 to 5, or arrays thereof, as well as a kit based thereon are used for the prognosis of breast cancer relapse and/or the prediction of survival of a patient diagnosed with breast cancer, preferably under endocrine treatment since surgical removal of the tumour.

Moreover, an additional aspect of the present invention is a kit comprising, for example: a bisulfite-containing reagent as well as at least one oligonucleotide whose sequences in each case correspond, are complementary, or hybridise under stringent or highly stringent conditions to a 18-base long segment of the sequences SEQ ID NOS: 1 to 5. Said kit may further comprise instructions for carrying out and evaluating the described method. In a further preferred embodiment, said kit may further comprise standard reagents for performing a CpG position-specific methylation analysis, wherein said analysis comprises one or more of the following techniques: MS-SNuPE, MSP, MethyLight™, HeavyMethyl™, COBRA, and nucleic acid sequencing. However, a kit along the lines of the present invention can also contain only part of the aforementioned components.

Typical reagents (*e.g.*, as might be found in a typical COBRA-based kit) for COBRA analysis may include, but are not limited to: PCR primers for specific gene (or methylation-altered DNA sequence or CpG island); restriction enzyme and appropriate buffer; gene-hybridisation oligo; control hybridisation oligo; kinase labelling kit for oligo probe; and radioactive nucleotides. Additionally, bisulfite conversion reagents may include: DNA denaturation buffer; sul-

fonation buffer; DNA recovery reagents or kits (*e.g.*, precipitation, ultrafiltration, affinity column); desulfonation buffer; and DNA recovery components.

Typical reagents (*e.g.*, as might be found in a typical MethyLight®-based kit) for MethyLight® analysis may include, but are not limited to: PCR primers for specific gene (or methylation-altered DNA sequence or CpG island); TaqMan® probes; optimised PCR buffers and deoxynucleotides; and Taq polymerase.

Typical reagents (*e.g.*, as might be found in a typical Ms-SNuPE-based kit) for Ms-SNuPE analysis may include, but are not limited to: PCR primers for specific gene (or methylation-altered DNA sequence or CpG island); optimised PCR buffers and deoxynucleotides; gel extraction kit; positive control primers; Ms-SNuPE primers for specific gene; reaction buffer (for the Ms-SNuPE reaction); and radioactive nucleotides. Additionally, bisulfite conversion reagents may include: DNA denaturation buffer; sulfonation buffer; DNA recovery reagents or kit (*e.g.*, precipitation, ultrafiltration, affinity column); desulfonation buffer; and DNA recovery components.

Typical reagents (*e.g.*, as might be found in a typical MSP-based kit) for MSP analysis may include, but are not limited to: methylated and unmethylated PCR primers for specific gene (or methylation-altered DNA sequence or CpG island), optimised PCR buffers and deoxynucleotides, and specific probes.

Specifically, the present invention is related to a method for characterising a cell proliferative disorder of the breast tissues and/or predicting the survival of a patient diagnosed with said disorder, comprising the steps of: a) detecting the expression of a nucleic acid or a polypeptide expressed from the PITX2 gene in an isolated biological sample representative of said cell proliferative disorders of the breast tissues and b) predicting therefrom the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient. Preferably, the method according to the present invention further comprises c) determining a suitable treatment regimen for the subject.

Preferred is a method according to the present invention, wherein the patient is characterised by being subject to adjuvant endocrine therapy comprising one or more treatments which target the estrogen receptor pathway or are involved in estrogen metabolism, production or se-

cretion.

Preferred is also a method according to the present invention, wherein said breast cell proliferative disorders are taken from the group comprising ductal carcinoma *in situ*, invasive ductal carcinoma, invasive lobular carcinoma, lobular carcinoma *in situ*, comedocarcinoma, inflammatory carcinoma, mucinous carcinoma, scirrhous carcinoma, colloid carcinoma, tubular carcinoma, medullary carcinoma, metaplastic carcinoma, and papillary carcinoma and papillary carcinoma *in situ*, undifferentiated or anaplastic carcinoma and Paget's disease of the breast.

According to another aspect of the method according to the present invention, said method is characterised in that the detection is carried out by a) contacting said biological sample with an antibody immunoreactive with the PITX2 polypeptide to form an immunocomplex; b) detecting said immunocomplex; and c) predicting therefrom the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient.

According to another aspect of the method according to the present invention, said method is characterised in that the detection is carried out by a) contacting said biological sample with an antibody immunoreactive with the PITX2 polypeptide to form an immunocomplex; b) detecting said immunocomplex; c) therefrom predicting the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient, and d) comparing the quantity of said immunocomplex to the quantity of immunocomplex formed under identical conditions with the same antibody and a control sample from one or more patients with a known prognosis.

According to yet another aspect of the method according to the present invention, the detection is carried out by a) contacting said biological sample with an antibody immunoreactive with the PITX2 polypeptide to form an immunocomplex; b) detecting said immunocomplex; and c) predicting therefrom the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient, and wherein an increase in quantity of said immunocomplex in the sample from said subject relative to said control sample is indicative of a bad prognosis.

Preferred is a method according to the present invention, wherein said immunoassay is a radioimmunoassay or an ELISA or a Western blot. Preferred is a method according to the present invention, wherein said detection is afforded by mRNA expression analysis. Most preferred is a method according to the present invention, comprising detecting the level of mRNA encoding a PITX2 polypeptide in a biological sample from a patient,

Preferred is a method according to the present invention, wherein a increased concentration of said mRNA above the concentration determined for an individual known to have a good prognosis indicates a bad prognosis.

According to yet another aspect of the method according to the present invention said method comprises the steps of: a) providing a polynucleotide probe which specifically hybridises or is identical to a polynucleotide consisting of SEQ ID NO: 19 or SEQ ID NO: 1, b) incubating said sample with said polynucleotide probe under high stringency conditions to form a specific hybridisation complex between a nucleic acid and said probe; and c) detecting said hybridisation complex.

Preferred is a method according to the present invention wherein said nucleic acid is mRNA or a cDNA derived therefrom.

Preferred is a method according to the present invention wherein the detecting step further comprises the steps of: a) producing a cDNA from mRNA in the sample; b) providing two oligonucleotides which specifically hybridise to regions flanking a segment of the cDNA; c) performing a polymerase chain reaction on the cDNA of step a) using the oligonucleotides of step b) as primers to amplify the cDNA segment; and d) detecting the amplified cDNA segment.

Yet another aspect of the present invention relates to a use of a polypeptide expressed from the PITX2 gene for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis. Preferably, said polypeptide is expressed from the PITX2 gene and used for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.

Preferred is a method according to the present invention wherein said detection comprises determining the genetic parameters of the gene PITX2, its promoter and/or regulatory elements. More preferred is a method according to the present invention wherein said detection comprises determining the epigenetic parameters of the gene PITX2, its promoter and/or regulatory elements.

Preferred is a method according to the present invention, wherein said detection comprises determining the methylation status of one or more CpG positions of a target nucleic acid within the gene PITX2, its promoter and/or regulatory elements, in particular through the methylation analysis of a genomic DNA sequence according to SEQ ID NO: 1. Preferred is further a method according to the present invention, wherein said detection comprises determining the methylation status of one or more CpG positions of a target nucleic acid characterised as being identical to or hybridising under stringent or moderately stringent conditions to a sequence out of the group of SEQ ID NOs 13, 18 and 19. Preferred is further a method according to the present invention, wherein the methylation analysis is afforded by contacting said target nucleic acid with one or more agents that convert cytosine bases that are unmethylated at the 5'-position thereof to a base that is detectably dissimilar to cytosine in terms of hybridisation properties.

Preferred is a method according to the present invention, wherein contacting said target nucleic acids with one or more agents comprises use of a solution selected from the group consisting of bisulfite, hydrogen sulfite, disulfite, and combinations thereof.

Yet another aspect of the present invention relates to the use of a set of oligomer probes comprising at least two oligomers, in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for detecting the cytosine methylation state and/or single nucleotide polymorphisms (SNPs) within one of the sequences according to SEQ ID NO: 1, and sequences complementary thereto.

Yet another aspect of the present invention relates to a method for manufacturing an arrangement of different oligomers (array) fixed to a carrier material for analysing diseases associated

with the methylation state of the CpG dinucleotides of one of SEQ ID NO: 1, and sequences complementary thereto wherein at least one oligomer, in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, is coupled to a solid phase.

Yet another aspect of the present invention relates to a composition of matter comprising the following: a) a nucleic acid comprising a sequence at least 18 bases in length of a segment of the chemically pretreated genomic DNA according to one of the sequences taken from the group comprising SEQ ID NO: 1 to SEQ ID NO: 5 and sequences complementary thereto, and b) a buffer comprising at least one of the following substances: 1 to 5 mM magnesium chloride, 100-500 μ M dNTP, 0.5-5 units/10ul of taq polymerase, an oligomer, in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto.

Preferably, the gene PITX2, its promoter and/or regulatory elements is used for predicting the survival of patients diagnosed with a cell proliferative disease. Preferred is the use of the mRNA of the gene PITX2 for predicting the survival of patients diagnosed with a cell proliferative disease.

Yet another aspect of the present invention relates to a method for predicting the survival of patients diagnosed with a cell proliferative disease according to the present invention, comprising: a) isolating or enriching genomic DNA from said biological sample ; b) treating the genomic DNA, or a fragment thereof, with one or more reagents to convert 5-position unmethylated cytosine bases to uracil or to another base that is detectably dissimilar to cytosine in terms of hybridisation properties; c) contacting the treated genomic DNA, or the treated fragment thereof, with an amplification enzyme and at least two primers comprising, in each case a contiguous sequence at least 18 nucleotides in length that is complementary to, or hybridises under moderately stringent or stringent conditions to a sequence selected from the

group consisting of SEQ ID NOS: 2 to 5, and complements thereof, wherein the treated DNA or a fragment thereof is either amplified to produce one or more amplicates, or is not amplified; and d) determining, based on the presence or absence of, or on the quantity or on a property of said amplicate, the methylation state of at least one CpG dinucleotide sequence of SEQ ID NO: 1, or an average, or a value reflecting an average methylation state of a plurality of CpG dinucleotide sequences of SEQ ID NO: 1.

Yet another aspect of the present invention relates to a method for detecting the survival of patients diagnosed with a cell proliferative disease of the breast according to the present invention, comprising the following steps of a) obtaining, from a subject, a biological sample having subject genomic DNA; b) treating the genomic DNA, or a fragment thereof, with one or more reagents to convert 5-position unmethylated cytosine bases to uracil or to another base that is detectably dissimilar to cytosine in terms of hybridisation properties; c) amplifying one or more fragments of the treated DNA such that only DNA originating from breast or breast cell proliferative disorder cells are amplified and d) detecting the amplicates or characteristics thereof and thereby deducing on the survival of patients diagnosed with a cell proliferative disease of the breast.

Yet another aspect of the present invention relates to a use of an oligomer, an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to an artificially modified, chemically pretreated DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis.

Yet another aspect of the present invention relates to a use of a nucleic acid comprising a sequence of at least 18 bases in length of a segment of the artificially modified, chemically pretreated, DNA according to one of the sequences taken from the group comprising SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis. Yet another aspect of the present invention relates to a use of an oligomer, an oligonucleotide or peptide nucleic acid (PNA)-oligomer,

said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to an artificially modified, chemically pretreated DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.

Yet another aspect of the present invention relates to a use of a nucleic acid comprising a sequence of at least 18 bases in length of a segment of the artificially modified, chemically pretreated, DNA according to one of the sequences taken from the group comprising SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast. Yet another aspect of the present invention relates to a use of a nucleic acid represented by their SEQ ID NO out of the group of nucleic acids according to SEQ ID NOS : 6, 7, 8, 9, 10, 14, 15, 22, 23, 24, 25, 26, 27, 28, 29, 30 and 31, for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis. Yet another aspect of the present invention relates to a use of a nucleic acid represented by their SEQ ID NO out of the group of nucleic acids according to SEQ ID NOS : 6, 7, 8, 9, 10, 14, 15, 22, 23, 24, 25, 26, 27, 28, 29, 30 and 31, for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.

In the context of this invention the terms "obtaining a biological sample" or "obtaining a sample from a subject", is supposed to comprise several different sources of such a sample, but always excludes the active retrieval of a sample from an individual patient, such as the performance of a biopsy. Included are the following examples: obtaining a sample, which was prior to the obtaining step taken from a patient in a biopsy or surgery, from a sample provider; obtaining a sample from a clinician, a surgeon or other medical personnel; obtaining a sample from a courier, who is bringing the sample from the clinician or practitioner or patient himself, for example to the analytic service station, as well as obtaining a sample such as a body fluid sample per post or from the hands of the patient himself. This is not meant to be a limiting list, but shall illustrate that it is not a feature of the invention that it needs to be carried out on the patient itself.

The term "biological material" relates to any material that is derived from a source, in particular an animal and/or human source, that contains or is suspected to contain genomic DNA.

One example of a biological material according to the present invention will be a biological sample.

In the context of the present invention, the term "CpG island" refers to a contiguous region of genomic DNA that satisfies the criteria of (1) having a frequency of CpG dinucleotides corresponding to an "Observed/Expected Ratio" >0.6, and (2) having a "GC Content" >0.5. CpG islands are typically, but not always, between about 0.2 to about 1 kb in length.

In the context of the present invention the term "regulatory region" of a gene is taken to mean nucleotide sequences which affect the expression of a gene. Said regulatory regions may be located within, proximal or distal to said gene. Said regulatory regions include but are not limited to constitutive promoters, tissue-specific promoters, developmental-specific promoters, inducible promoters and the like. Promoter regulatory elements may also include certain enhancer sequence elements that control transcriptional or translational efficiency of the gene.

In the context of the present invention, the term "methylation" refers to the presence or absence of 5-methylcytosine ("5-mCyt") at one or a plurality of CpG dinucleotides within a DNA sequence.

In the context of the present invention the term "methylation state" is taken to mean the degree of methylation present in a nucleic acid of interest, this may be expressed in absolute or relative terms i.e. as a percentage or other numerical value or by comparison to another tissue and therein described as hypermethylated, hypomethylated or as having significantly similar or identical methylation status.

In the context of the present invention, the term "hemi-methylation" or "hemimethylation" refers to the methylation state of a palindromic CpG methylation site, where only a single cytosine in one of the two CpG dinucleotide sequences of the double stranded CpG methylation site is methylated (*e.g.*, 5'-NNC^MGNN-3' (top strand): 3'-NNGCNN-5' (bottom strand)).

In the context of the present invention, the term "hypermethylation" refers to the average methylation state corresponding to an *increased* presence of 5-mCyt at one or a plurality of

CpG dinucleotides within a DNA sequence of a test DNA sample, relative to the amount of 5-mCyt found at corresponding CpG dinucleotides within a normal control DNA sample.

In the context of the present invention, the term "hypomethylation" refers to the average methylation state corresponding to a *decreased* presence of 5-mCyt at one or a plurality of CpG dinucleotides within a DNA sequence of a test DNA sample, relative to the amount of 5-mCyt found at corresponding CpG dinucleotides within a normal control DNA sample.

In the context of the present invention, the term "microarray" refers broadly to both "DNA microarrays," and 'DNA chip(s),' as recognised in the art, encompasses all art-recognised solid supports, and encompasses all methods for affixing nucleic acid molecules thereto or synthesis of nucleic acids thereon.

"Genetic parameters" are mutations and polymorphisms of genes and sequences further required for their regulation. To be designated as mutations are, in particular, insertions, deletions, point mutations, inversions and polymorphisms and, particularly preferred, SNPs (single nucleotide polymorphisms).

"Epigenetic modifications" or "epigenetic parameters" are modifications of DNA bases of genomic DNA and sequences further required for their regulation, in particular, cytosine methylations thereof. Further epigenetic parameters include, for example, the acetylation of histones which, however, cannot be directly analysed using the described method but which, in turn, correlate with the DNA methylation.

In the context of the present invention, the term "bisulfite reagent" refers to a reagent comprising bisulfite, disulfite, hydrogen sulfite or combinations thereof, useful as disclosed herein to distinguish between methylated and unmethylated CpG dinucleotide sequences.

In the context of the present invention, the term "Methylation assay" refers to any assay for determining the methylation state of one or more CpG dinucleotide sequences within a sequence of DNA.

In the context of the present invention, the term "MS.AP-PCR" (Methylation-Sensitive Arbitrarily-Primed Polymerase Chain Reaction) refers to the art-recognised technology that allows

for a global scan of the genome using CG-rich primers to focus on the regions most likely to contain CpG dinucleotides, and described by Gonzalgo et al., *Cancer Research* 57:594-599, 1997.

In the context of the present invention, the term "MethyLight" refers to the art-recognised fluorescence-based real-time PCR technique described by Eads et al., *Cancer Res.* 59:2302-2306, 1999.

In the context of the present invention, the term "HeavyMethyl™" assay, in the embodiment thereof implemented herein, refers to a HeavyMethyl™ MethyLight assay, which is a variation of the MethyLight assay, wherein the MethyLight assay is combined with methylation specific *blocking* probes covering CpG positions between the amplification primers.

The term "Ms-SNuPE" (Methylation-sensitive Single Nucleotide Primer Extension) refers to the art-recognised assay described by Gonzalgo & Jones, *Nucleic Acids Res.* 25:2529-2531, 1997.

In the context of the present invention the term "MSP" (Methylation-specific PCR) refers to the art-recognised methylation assay described by Herman et al. *Proc. Natl. Acad. Sci. USA* 93:9821-9826, 1996, and by US Patent No. 5,786,146.

In the context of the present invention the term "COBRA" (Combined Bisulfite Restriction Analysis) refers to the art-recognised methylation assay described by Xiong & Laird, *Nucleic Acids Res.* 25:2532-2534, 1997.

In the context of the present invention the term "hybridisation" is to be understood as a bond of an oligonucleotide to a complementary sequence along the lines of the Watson-Crick base pairings in the sample DNA, forming a duplex structure.

"Stringent hybridisation conditions," as defined herein, involve hybridising at 68°C in 5x SSC/5x Denhardt's solution/1.0% SDS, and washing in 0.2x SSC/0.1% SDS at room temperature, or involve the art-recognised equivalent thereof (e.g., conditions in which a hybridisation is carried out at 60°C in 2.5 x SSC buffer, followed by several washing steps at 37°C in a low buffer concentration, and remains stable). Moderately stringent conditions, as defined

herein, involve including washing in 3x SSC at 42°C, or the art-recognised equivalent thereof. The parameters of salt concentration and temperature can be varied to achieve the optimal level of identity between the probe and the target nucleic acid. Guidance regarding such conditions is available in the art, for example, by Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, N.Y.; and Ausubel et al. (eds.), 1995, *Current Protocols in Molecular Biology*, (John Wiley & Sons, N.Y.) at Unit 2.10.

"Background DNA" as used herein refers to any nucleic acids which originate from sources other than colon cells.

In the context of this application "survival" is meant to describe the time from diagnosis or start of treatment to an endpoint, which may be either the time of death (considering any reason for death or only death from breast cancer), or the time of recurrence of breast cancer (for example in form of metastases), which may be local or distant, or the time of occurrence of any breast cancer associated disease. Therefore "predicting the survival" is meant to comprise predicting the disease free survival, as well as the overall survival or any other consideration of time between diagnosis and endpoint of treatment. However, as it is obvious in the state of the art, a precise prediction of life time is generally impossible whether it is based on a biomarker analysis, or any other prognostic tools, it is understood throughout the invention that said term "prediction of survival" or "predicting the survival" is used to describe the risk of patient to suffer from a recurrence of metastasis or other disease caused by the original breast cell proliferative disease the patient was diagnosed with (also termed "risk of relapse"). Said risk can be predicted with a certain probability or likelihood. It is also clear, that predicting the survival is meant to comprise the determination of the likelihood or probability whether a subject or patient will survive for a longer or shorter period of time.

Throughout this invention it is preferred that said survival is characterised as the disease free or the overall survival. It is especially preferred that survival is understood as disease free survival. Disease free survival is understood as absence of recurrence of cancer (local or distant).

The terms "endocrine therapy" or "endocrine treatment" is meant to comprise any therapy, treatment or treatments targeting the estrogen receptor pathway or estrogen synthesis pathway or estrogen conversion pathway, which is involved in estrogen metabolism, production or

secretion. Said treatments include, but are not limited to estrogen receptor modulators, estrogen receptor down-regulators, aromatase inhibitors, ovarian ablation, LHRH analogues and other centrally acting drugs influencing estrogen production.

The term "monotherapy" is used to explain that no other treatment is given in addition or to support said monotherapy.

In the context of the present invention the term "chemotherapy" is taken to mean the use of drugs or chemical substances to treat cancer. This definition excludes radiation therapy (treatment with high energy rays or particles), hormone therapy (treatment with hormones or hormone analogues (synthetic substitutes) and surgical treatment.

In the context of the present invention the term "adjuvant treatment" is taken to mean a therapy of a cancer patient immediately following an initial non chemotherapeutical therapy, e.g. surgery. In general, the purpose of an adjuvant therapy is to provide a significantly smaller risk of recurrences compared without the adjuvant therapy.

In the context of the present invention the term "determining a suitable treatment regimen for the subject" is taken to mean a treatment regimen (i.e. a single therapy or a combination of different therapies that are used for the prevention and/or treatment of the cancer in the patient) for the cancer patient that is started, modified and/or ended based or essentially based or at least partially based on the results of the analysis according to the present invention. One example is starting an adjuvant endocrine therapy after surgery, another would be to modify the dosage of a particular chemotherapy. The determination can, in addition to the results of the analysis according to the present invention, be based on personal characteristics of the subject to be treated. In most cases, the actual determination of the suitable treatment regimen for the subject will be performed by the attending physician or doctor.

While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples and figures serve only to illustrate the invention and is not intended to limit the invention within the principles and scope of the broadest interpretations and equivalent configurations thereof.

In the accompanying sequence protocol and the figures,

SEQ ID NO: 1 shows the nucleic acid sequence of the human gene PITX2,

SEQ ID NOS: 2 to 5 show chemically pretreated nucleic acid sequences of the gene PITX2, according to table 1.

SEQ ID NOS: 6 to 9 show the nucleic acid sequences of those primers and probes useful to predict the survival of breast cancer patients according to the invention as described in example 4.

SEQ ID NOS: 14 to 17 show the nucleic acid sequences of those primers and probes useful to predict the survival of breast cancer patients according to the invention as described in example 5.

SEQ ID NOS: 10 to 12 show the nucleic acid sequences of primers and probes according to a control gene used in the example 4 and 5.

SEQ ID NO: 13 shows a subsequence of SEQ ID 1, which represents the nucleic acid sequence of the human gene PITX2.

SEQ ID NO: 18 shows an amino acid sequence of the polypeptide encoded by the gene PITX2. The amino acid sequence of the polypeptide encoded by the gene PITX2 is also illustrated in figure 10.

FIGURES

Figure 1 presents a scheme to illustrate a preferred application of the method according to the invention. Along the Y axis tumour(s) mass (or size) increases, wherein the line '3' indicates the limit of detectability of said tumour mass. The X axis represents time (such as in life time of a patient). Accordingly said figure illustrates a simplified model of an Stage 1-3 breast tumour wherein primary treatment was surgery (at point 1), followed by adjuvant therapy with Tamoxifen, as an example for an endocrine treatment. In a first scenario a patient without relapse during endocrine treatment (4) is shown as remaining below the limit of detectability for the duration of the observation. A patient with relapse of the cancer (5) has a period of disease free survival (2) followed by relapse when the carcinoma mass reaches the level of detectability.

Figure 2 shows the result of the assay (QM assay) as described in example 4: A Kaplan-Meier estimated metastasis-free survival curve for 3 CpG sites of the PITX2 gene by means of Real-Time methylation specific probe analysis (QM assay). The lower curve shows the proportion of metastasis free patients in the population with above median methylation levels, the upper curve shows the proportion of metastasis free patients in the population with below median

methylation levels. The X axis shows the metastasis free survival times of the patients in months, and the Y axis shows the proportion of metastasis free survival patients.

Figure 3 shows the result of the chip hybridisation experiment as described in example 2. A Kaplan-Meier estimated metastasis-free survival curves for 2 CpG positions of the PITX2 gene by means of methylation specific detection oligo hybridisation analysis. The lower curve shows the proportion of metastasis free patients in the population with above median methylation levels, the upper curve shows the proportion of metastasis free patients in the population with below median methylation levels. The X axis shows the metastasis free survival times of the patients in months, and the Y axis shows the proportion of metastasis free survival patients.

Figure 4 shows the Kaplan-Meier estimated metastasis-free survival curves for 2 CpG positions of the PITX2 gene by means of methylation specific detection oligo hybridisation analysis. The lower line shows the proportion of metastasis free patients in the population of 55 patients with above median methylation levels, the upper curve shows the proportion of metastasis free patients in the population of 54 patients with below median methylation levels. The X axis shows the metastasis free survival times of the patients in years, and the Y axis shows the proportion of metastasis free survival patients in %. This resulted from a first data set that was achieved in a first study.

Figure 5 shows the Kaplan-Meier estimated metastasis-free survival curves for 6 different CpG positions located within the preferred region of the PITX2 gene (SEQ ID 13) by means of methylation specific detection oligo hybridisation analysis. The lower line shows the proportion of metastasis free patients in the population of 118 patients with above median methylation levels, the upper curve shows the proportion of metastasis free patients in the population of 118 patients with below median methylation levels. The X axis shows the metastasis free survival times of the patients in years, and the Y axis shows the proportion of metastasis free survival patients in %. This resulted from a second data set that was achieved in a second study.

Figure 6 shows the Kaplan-Meier estimated metastasis-free survival curves for 6 different CpG positions located within the preferred region of the PITX2 gene (SEQ ID 13) by means of methylation specific detection oligo hybridisation analysis. This time only a subpopulation

of 148 patients, characterised by a tumour at grade G1 or G2, was analysed: The lower curve shows the proportion of metastasis free patients in the population of 74 patients with above median methylation levels, the upper curve shows the proportion of metastasis free patients in the population of 74 patients with below median methylation levels. The X axis shows the metastasis free survival times of the patients in years, and the Y axis shows the proportion of metastasis free survival patients in %. This resulted from a second data set that was achieved in the second example.

Figure 7 shows the Kaplan-Meier estimated metastasis-free survival curves for 4 different CpG positions located within the preferred region of the PITX2 gene (SEQ ID 13) by means of methylation specific detection oligo hybridisation analysis. This time a subpopulation of 224 patients, characterised by a tumour of stage 1 or 2 (T1 or T2), was analysed: The lower curve shows the proportion of metastasis free patients in the population of 112 patients with above median methylation levels, the upper curve shows the proportion of metastasis free patients in the population of 112 patients with below median methylation levels. The X axis shows the metastasis free survival times of the patients in years, and the Y axis shows the proportion of metastasis free survival patients in %. This resulted from the second data set that was achieved in the second example.

Figure 8 shows the disease-free survival curves for a combination of two oligonucleotides each from the genes TBC1D3 and CDK6, and one oligonucleotide from the gene PITX2 covering two CpG sites. The black curve shows the proportion of disease free patients in the population with above median methylation scores, the grey curve shows the proportion of disease free patients in the population with below median methylation scores.

Figure 9 shows the plot according to Figure 8 and the classification of the sample set by means of the St. Gallen method. The unbroken lines represent the methylation analysis wherein the black curve shows the proportion of disease free patients in the population with above median methylation scores, the grey curve shows the proportion of disease free patients in the population with below median methylation scores. The broken lines represent the St. Gallen classification of the sample set wherein the black curve shows the disease free survival time of the high risk group and the grey curve shows the disease free survival of the low risk group.

Figure 10 illustrates the amino acid sequence of the polypeptide encoded by the gene PITX2.

EXAMPLES

EXAMPLE 1 : Study 1

The first study was based on a population of 109 patients, comprising patients of both nodal statuses N0 and N+. All patients were ER+ (estrogen receptor positive). All patients received Tamoxifen monotherapy immediately after surgery or diagnosis. The samples were analysed using Epigenomics' chip technology with two chip panels representing altogether 117 candidate genes. For further details see patent application WO 04/035803 and EP 03 090 432.0, which is hereby incorporated by reference. In this study one significant marker gene was found. The methylation status of PITX2, coding for a transcription factor, was correlated statistically significant with disease-free survival under adjuvant Tamoxifen treatment. A Cox regression model that includes the nodal status of the patient at the time of diagnosis was applied.

The result from this study – with respect to PITX2 - is illustrated in Figure 4. The X axis shows the metastasis free survival times of the patients in years, and the Y axis shows the proportion of metastasis free survival patients in %. Amongst the 54 patients (upper line) with below median methylation levels a higher percentage has a significantly longer metastasis free survival time, than amongst the 55 patients with above median methylation levels (lower line). To illustrate the result: At time of 10 years after surgery under tamoxifen monotherapy, more than 75% of the patients with low methylation in PITX2 are still metastasis free, whereas less than 60% of the patients with high methylation in PITX2.

As the survival of a breast cancer patient is known to also be correlated to the patient's nodal status, the differentiating power of the marker in this mixed population is expected to be less than in a homogenous population.

Another study was performed to analyse whether the same marker can be identified independently, in a completely different set of patient samples and also to characterise the differential power towards predicting survival for a sub-group of patients, all being N0.

EXAMPLE 2 : Study 2

The second study was based on samples from 236 patients from 5 different centres, wherein all patients were N0 (nodal status negativ), and older than 35 years. In all cases the surgery was performed before 1998. All patients were ER+ (estrogen receptor positive), and the tumours were graded to be T1-3, G1-3. In this study all patients received Tamoxifen directly after surgery, and the outcome was assessed as the length of disease-free survival. In order to be as representative as possible for the final target group, the patients and their tumor samples had to fulfil the following criteria:

The range and median follow-up of patients were the following:

Median: 64.5 months

Range: 3 months to 142 months

(calculated based on patients who were disease-free at end of observation time).

Analysis of the methylation patterns of patient samples treated with Tamoxifen as an adjuvant therapy immediately following surgery (see Figure 1) is shown in the plots according to Figures 5 to 7. For the amplicate, the mean methylation over 4 oligo-pairs for that amplicate was calculated and the population split into groups according to their mean methylation values, wherein one group was composed of individuals with a methylation score higher than the median and a second group composed of individuals with a methylation score lower than the median.

The primer oligonucleotides used to generate the amplicate, that was analysed in the array experiment were these :

Array Primer PITX2_Q21: GTAGGGGAGGGAAGTAGATGT (SEQ ID 22)

Array Primer PITX2_R23: TCCTCAACTCTACAAACCTAAAA (SEQ ID 23)

The according genomic region of said amplicate is given in SEQ ID 13.

The sequences of the oligonucleotides used in this array experiment were the following:

SEQ ID NO 24 : AGTCGGGAGAGCGAAA

SEQ ID NO 25 : AGTTGGGAGAGTGAAA

SEQ ID NO 26 : AAGAGTCGGGAGTCGGA

SEQ ID NO 27: _AAGAGTTGGGAGTTGGA

SEQ ID NO 28: GGTCGAAGAGTCGGGA

SEQ ID NO 29: GGTTGAAGAGTTGGGA

SEQ ID NO 30: ATGTTAGCGGGTCGAA

SEQ ID NO 31: TAGTGGGTTGAAGAGT

When the data derived from analysing 6 different CpG sites, located within the preferred amplified region of the PITX2 gene by means of methylation specific detection oligo hybridisation analysis were plotted as Kaplan-Meier estimated metastasis-free survival curves, it can be seen that the differential power of the marker PITX2 increased with selecting for N0 patients. This is shown in figures 5 to 7. The X axis shows the metastasis free survival times of the patients in years, and the Y axis shows the proportion of metastasis free survival patients in %. The lower curve shows the proportion of metastasis free patients in the population with above median methylation levels, and the upper curve shows the proportion of metastasis free patients in the population with below median methylation levels.

For example, as illustrated in figure 5, 10 years after surgery only about 65% of the patients of the 118 patients with the higher methylation status are metastasis free, whereas about 90% of the 118 patients with lower methylation status are metastasis free.

As illustrated in figure 6 when looking at the analogous Kaplan-Meier analysis for a subpopulation of 148 patients, characterized by a tumour at stage G1 or G2 this differential power increases again: 10 years after surgery only about 60% of the 74 patients with the higher methylation status are metastasis free, whereas about 95% of the 74 patients with lower methylation status are metastasis free.

Figure 7 illustrates how the survival is also correlated to the tumour stage at surgery by showing the analogous Kaplan-Meier analysis for a subpopulation of 150 patients, characterised by a tumour stage of T1 or T2: The number of patients with 10 years MFS is about 68% of patients of the 112 with the higher methylation status, whereas about 95% of the 112 patients with lower methylation status are metastasis free.

EXAMPLE 3:

The accuracy of the differentiation between the different groups was further increased by combining multiple oligonucleotides from different genes. As described in the text it was recognised that adding additional informative markers to the analysis could potentially increase the prognostic power of a survival test. Therefore it was calculated how a combination of two methylation specific oligonucleotides each from the genes TBC1D3 and CDK6, and one oli-

gonucleotide from the gene PITX2 would differentiate the groups of good or bad prognosis. The result is shown in figure 8 as the according Kaplan-Meier curve.

Figure 9 shows -on top of Figure 8- the classification of the patients from the sample set by means of the St.Gallen method (the current method of choice for estimating disease free survival), thereby showing the improved effectiveness of methylation analysis over current methods, in particular post 80 months.

EXAMPLE 4: Real time Quantitative methylation analysis

Genomic DNA was analysed using the Real Time PCR technique after bisulfite conversion. In this analysis four oligonucleotides were used in each reaction. Two non methylation specific PCR primers were used to amplify a segment of the treated genomic DNA containing a methylation variable oligonucleotide probe binding site. Two oligonucleotide probes competitively hybridise to the binding site, one specific for the methylated version of the binding site, the other specific to the unmethylated version of the binding site. Accordingly, one of the probes comprises a CpG at the methylation variable position (i.e. anneals to methylated bisulphite treated sites) and the other comprises a TpG at said position (i.e. anneals to unmethylated bisulphite treated sites). Each species of probe is labelled with a 5' fluorescent reporter dye and a 3' quencher dye wherein the CpG and TpG oligonucleotides are labelled with different dyes.

The reactions are calibrated by reference to DNA standards of known methylation levels in order to quantify the levels of methylation within the sample. The DNA standards were composed of bisulfite treated phi29 amplified genomic DNA (i.e. unmethylated), and/or phi29 amplified genomic DNA treated with Sss1 methylase enzyme (thereby methylating each CpG position in the sample), which is then treated with bisulfite solution. Seven different reference standards were used with 0%, (i.e. phi29 amplified genomic DNA only), 5%, 10%, 25%, 50%, 75% and 100% (i.e. phi29 Sss1 treated genomic only).

The amount of sample DNA amplified is quantified by reference to the gene (β -actin (ACTB)) to normalise for input DNA. For standardisation the primers and the probe for analysis of the ACTB gene lack CpG dinucleotides so that amplification is possible regardless of methylation levels. As there are no methylation variable positions, only one probe oligonucleotide is required.

The following oligonucleotides were used in the reaction to amplify the control amplificate:

Control Primer1: TGGTGATGGAGGAGGTTTAGTAAGT (SEQ ID NO: 10)

Control Primer2: AACCAATAAAACCTACTCCTCCCTTAA (SEQ ID NO: 11)

Control Probe: 6FAM-ACCACCACCCAACACACAATAACAAACACA-TAMRA or Dabcyl (SEQ ID NO: 12)

The nucleic acid sequence of the gene PITX2 is given in (SEQ ID 1), after treatment with bisulfite two different strands are generated, and each of the strands is represented twice, once in a prior to treatment methylated version (SEQ ID 2 and 3) and once in the prior to treatment unmethylated form (SEQ ID 4 and 5), which are characterised as containing no cytosine bases (despite of those 5' adjacent to a guanine and methylated before treatment).

The following primers are used to generate an amplificate within the PITX2 sequence comprising the CpG sites of interest:

Primers for PITX bisulfite amplificate length : 144 bp

PITX2R02: GTAGGGGAGGGAAGTAGATGTT (SEQ ID NO: 6)

PITX2Q02: TTCTAATCCTCCTTCCACAATAA (SEQ ID NO: 7)

The genomic region according to the generated amplificate of 144 bp in length is given in SEQ ID NO 18.

Probes:

PITX2cg1: FAM-AGTCGGAGTCGGGAGAGCGA-Darquencher (SEQ ID NO: 8)

As an alternative quencher TAMRA was also used in additional experiments:

FAM-AGTCGGAGTCGGGAGAGCGA-TAMRA

PITX2tg1: YAKIMA YELLOW-AGTTGGAGTTGGGAGAGTGAAAGGAGA-Darquencher (SEQ ID NO: 9)

In additional experiments we also used :

VIC- AGTTGGAGTTGGGAGAGTGAAAGGAGA -TAMRA

The extent of methylation at a specific locus was determined by the following formula:

methylation rate= $100 * I(CG) / (I(CG) + I(TG))$

(I = Intensity of the fluorescence of CG-probe or TG-probe)

PCR components were ordered from Eurogentec :

3 mM MgCl₂ buffer, 10x buffer, Hotstart TAQ

Program (45 cycles): 95 °C, 10 min; 95 °C, 15 sec; 62 °C, 1 min

This assay was performed on 236 samples identical to those used in Example 2. The result is shown in figure 2. Figure 2 shows the Kaplan-Meier estimated disease-free survival curves for 3 CpG positions of the PITX2 gene by means of Real-Time methylation specific probe analysis, as described above. The lower curve shows the proportion of disease free patients in the population with above median methylation levels, the upper curve shows the proportion of disease free patients in the population with below median methylation levels. The X axis shows the disease free survival times of the patients in months, and the Y- axis shows the proportion of disease free survival patients. The p-value (probability that the observed distribution occurred by chance) was calculated as 0.0031, thereby confirming the data obtained by means of array analysis.

For comparison, figure 3 illustrates the result from the array analysis of said gene, according to the chip hybridisation experiment described in Example 2, wherein detection oligos were used (for details see EP 03 090 432.0, which is incorporated by reference). The p-value (probability that the observed distribution occurred by chance) was calculated as 0.0011.

EXAMPLE 5

Another QM assay was developed in our hands, which also performed very well. The following PITX2 specific oligonucleotides were employed to generate an amplificate of 164 bp. The oligonucleotides are specific for three co-methylated CpG positions. :

Primers for PITX2 bisulfite amplificate with a length of 162 bp :

PITX2O2: AACATCTACTTCCCTCCCCTAC (SEQ ID NO: 14)

PITX2P3: GTTAGTAGAGATTTTATTAAATTTATTGTAT (SEQ ID NO: 15)

The genomic region according to the generated amplifacte of 162 bp in length is given in SEQ ID NO 19.

Probes (from ABI):

PITX2-IIcg1: FAM-TTCGGTTGCGCGGT-MGBNQF (SEQ ID NO: 16)

PITX2-IItg1: VIC-TTTGGTTGTGTGGTTG- MGBNQF (SEQ ID NO: 17)

The extent of methylation at a specific locus was determined by the following formula:

methylation rate = $100 * I(CG) / (I(CG) + I(TG))$

(I = Intensity of the fluorescence of CG-probe or TG-probe)

PCR components were ordered from Eurogentec : 2,5 mM MgCl₂ buffer, 10x buffer, Hotstart
TAQ

Program (45 cycles): 95 °C, 10 min; 95 °C, 15 sec; 60 °C, 1 min

05. Juni 2004

CLAIMS

1. A method for characterising a cell proliferative disorder of the breast tissues and/or predicting the survival of a patient diagnosed with said disorder, comprising the steps of:
 - (a) detecting the expression of a nucleic acid or a polypeptide expressed from the PITX2 gene in an isolated biological sample representative of said cell proliferative disorders of the breast tissues and
 - (b) therefrom predicting the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient.
2. The method according to claim 1 further comprising
 - (c) determining a suitable treatment regimen for the subject.
3. The method of claim 1, wherein said patient is characterised by being subject to adjuvant endocrine therapy comprising one or more treatments which target the estrogen receptor pathway or are involved in estrogen metabolism, production or secretion.
4. The method of claim 1, wherein said breast cell proliferative disorders are taken from the group comprising ductal carcinoma *in situ*, invasive ductal carcinoma, invasive lobular carcinoma, lobular carcinoma *in situ*, comedocarcinoma, inflammatory carcinoma, mucinous carcinoma, scirrhous carcinoma, colloid carcinoma, tubular carcinoma, medullary carcinoma, metaplastic carcinoma, and papillary carcinoma and papillary carcinoma *in situ*, undifferentiated or anaplastic carcinoma and Paget's disease of the breast.
5. The method according to claim 1 characterised in that the detection is carried out by
 - a) contacting said biological sample with an antibody immunoreactive with the PITX2 polypeptide to form an immunocomplex;
 - b) detecting said immunocomplex; and
 - c) predicting therefrom the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient.
6. The method according to claim 1 characterised in that the detection is carried out by

- a) contacting said biological sample with an antibody immunoreactive with the PITX2 polypeptide to form an immunocomplex;
- b) detecting said immunocomplex;
- c) therefrom predicting the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient, and
- d) comparing the quantity of said immunocomplex to the quantity of immunocomplex formed under identical conditions with the same antibody and a control sample from one or more patients with a known prognosis.

7. The method according to claim 1 characterised in that the detection is carried out by

- a) contacting said biological sample with an antibody immunoreactive with the PITX2 polypeptide to form an immunocomplex;
 - b) detecting said immunocomplex; and
 - c) therefrom predicting the survival of said patient, characteristics of said cell proliferative disorder, and/or prognosis of said patient,
- and wherein an increase in quantity of said immunocomplex in the sample from said subject relative to said control sample is indicative of a bad prognosis.

8. The method of claim 5, wherein said immunoassay is a radioimmunoassay or an ELISA or a Western blot.

9. The method of claim 1, wherein said detection is afforded by mRNA expression analysis.

10. The method of claim 9, comprising detecting the level of mRNA encoding a PITX2 polypeptide in a biological sample from a patient,

11. A method according to claim 10, wherein a increased concentration of said mRNA above the concentration determined for an individual known to have a good prognosis indicates a bad prognosis.

12. The method of claim 10, comprising the steps of:

- (a) providing a polynucleotide probe which specifically hybridises or is identical to a polynucleotide consisting of SEQ ID NO: 19 or SEQ ID NO: 1,

- (b) incubating said sample with said polynucleotide probe under high stringency conditions to form a specific hybridisation complex between a nucleic acid and said probe;
- (c) detecting said hybridisation complex.

13. The method according to claim 12 wherein said nucleic acid is mRNA or a cDNA derived therefrom.

14. The method according to claim 13 wherein the detecting step further comprises the steps of:

- a) producing a cDNA from mRNA in the sample;
- b) providing two oligonucleotides which specifically hybridise to regions flanking a segment of the cDNA;
- c) performing a polymerase chain reaction on the cDNA of step a) using the oligonucleotides of step b) as primers to amplify the cDNA segment; and
- d) detecting the amplified cDNA segment.

15. Use of a polypeptide expressed from the PITX2 gene for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis.

16. Use of a polypeptide expressed from the PITX2 gene for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.

17. The method of claim 1 wherein said detection comprises determining the genetic parameters of the gene PITX2, its promoter and/or regulatory elements.

18. The method of claim 1 wherein said detection comprises determining the epigenetic parameters of the gene PITX2, its promoter and/or regulatory elements.

19. The method of claim 1, wherein said detection comprises determining the methylation status of one or more CpG positions of a target nucleic acid within the gene PITX2, its promoter and/or regulatory elements, in particular through the methylation analysis of a genomic DNA sequence according to SEQ ID NO: 1.

20. The method of claim 1, wherein said detection comprises determining the methylation status of one or more CpG positions of a target nucleic acid characterised as being identical to or hybridising under stringent or moderately stringent conditions to a sequence out of the group of SEQ ID NOs 13, 18 and 19.
21. The method of claim 19, wherein the methylation analysis is afforded by contacting said target nucleic acid with one or more agents that convert cytosine bases that are unmethylated at the 5'-position thereof to a base that is detectably dissimilar to cytosine in terms of hybridisation properties.
22. The method of claim 21, wherein contacting said target nucleic acids with one or more agents comprises use of a solution selected from the group consisting of bisulfite, hydrogen sulfite, disulfite, and combinations thereof.
23. Use of a set of oligomer probes comprising at least two oligomers, in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for detecting the cytosine methylation state and/or single nucleotide polymorphisms (SNPs) within one of the sequences according to SEQ ID NO: 1, and sequences complementary thereto.
24. A method for manufacturing an arrangement of different oligomers (array) fixed to a carrier material for analysing diseases associated with the methylation state of the CpG dinucleotides of one of SEQ ID NO: 1, and sequences complementary thereto wherein at least one oligomer, in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, is coupled to a solid phase.
25. A composition of matter comprising the following:
- a nucleic acid comprising a sequence at least 18 bases in length of a segment of the

chemically pretreated genomic DNA according to one of the sequences taken from the group comprising SEQ ID NO: 1 to SEQ ID NO: 5 and sequences complementary thereto, and

- a buffer comprising at least one of the following substances: 1 to 5 mM Magnesium Chloride, 100-500 μ M dNTP, 0.5-5 units/10ul of taq polymerase, an oligomer, in particular an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to a pretreated genomic DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto.

26. Use of the gene PITX2, its promoter and/or regulatory elements for predicting the survival of patients diagnosed with a cell proliferative disease.

27. Use of the mRNA of the gene PITX2 for predicting the survival of patients diagnosed with a cell proliferative disease.

28. A method for predicting the survival of patients diagnosed with a cell proliferative disease according to claim 19, comprising:

- a) isolating or enriching genomic DNA from said biological sample ;
- b) treating the genomic DNA, or a fragment thereof, with one or more reagents to convert 5-position unmethylated cytosine bases to uracil or to another base that is detectably dissimilar to cytosine in terms of hybridisation properties;
- c) contacting the treated genomic DNA, or the treated fragment thereof, with an amplification enzyme and at least two primers comprising, in each case a contiguous sequence at least 18 nucleotides in length that is complementary to, or hybridises under moderately stringent or stringent conditions to a sequence selected from the group consisting of SEQ ID NOS: 2 to 5, and complements thereof, wherein the treated DNA or a fragment thereof is either amplified to produce one or more amplicates, or is not amplified; and
- d) determining, based on the presence or absence of, or on the quantity or on a property of said amplicate, the methylation state of at least one CpG dinucleotide sequence of SEQ ID NO: 1, or an average, or a value reflecting an average methylation state of a plurality of CpG dinucleotide sequences of SEQ ID NO: 1.

29. A method for detecting the survival of patients diagnosed with a cell proliferative disease of the breast according to claim 19, comprising the following steps of

- a) obtaining, from a subject, a biological sample having subject genomic DNA;
- b) treating the genomic DNA, or a fragment thereof, with one or more reagents to convert 5-position unmethylated cytosine bases to uracil or to another base that is detectably dissimilar to cytosine in terms of hybridisation properties;
- c) amplifying one or more fragments of the treated DNA such that only DNA originating from breast or breast cell proliferative disorder cells are amplified
- d) detecting the amplicates or characteristics thereof and thereby deducing on the survival of patients diagnosed with a cell proliferative disease of the breast.

30. Use of an oligomer, an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to an artificially modified, chemically pretreated DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis.

31. Use of a nucleic acid comprising a sequence of at least 18 bases in length of a segment of the artificially modified, chemically pretreated, DNA according to one of the sequences taken from the group comprising SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis.

32. Use of an oligomer, an oligonucleotide or peptide nucleic acid (PNA)-oligomer, said oligomer comprising in each case at least one base sequence having a length of at least 9 nucleotides which is complementary to, or hybridises under moderately stringent or stringent conditions to an artificially modified, chemically pretreated DNA according to one of the SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.

33. Use of a nucleic acid comprising a sequence of at least 18 bases in length of a segment of the artificially modified, chemically pretreated, DNA according to one of the sequences taken from the group comprising SEQ ID NO: 2 to SEQ ID NO: 5 and sequences complementary thereto, for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.

34. Use of a nucleic acid represented by their SEQ ID NO out of the group of nucleic acids according to SEQ ID NOS : 6, 7, 8, 9, 10, 14, 15, 22, 23, 24, 25, 26, 27, 28, 29, 30 and 31, for differentiating or distinguishing between patients diagnosed with breast cancer, who have a good survival prognosis and patients who have a bad survival prognosis.

35. Use of a nucleic acid represented by their SEQ ID NO out of the group of nucleic acids according to SEQ ID NOS : 6, 7, 8, 9, 10, 14, 15, 22, 23, 24, 25, 26, 27, 28, 29, 30 and 31, for prediction of survival of a patient diagnosed with a cell proliferative disorder of the breast.


Abstract

EPO - Munich
69

05. Juni 200

The present invention relates to methods for predicting the survival of a human being diagnosed with a cell proliferative disorder of the breast tissues, characterised by a step of determining the expression level of PITX2 or the genetic or the epigenetic modifications of the genomic DNA associated with the gene PITX2. The invention also relates to sequences, oligonucleotides and antibodies which can be used within the described methods.

Figure 1



EPO - Munich
69
0 5. Juni 2004

Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8





Figure 9



Sequence listing

EPO - Munich
69

05. Juni 2004

<110> Epigenomics AG

<120> PITX2 - a marker to predict survival of patients diagnosed with breast cell proliferative disease

<160> 12

<210> 1

<211> 9001

<212> DNA

<213> Homo Sapiens

<400> 1

agctgatgga	cttgctaaat	ttctttcttc	ttttttcttt	ttcatattat	ttgctagcca	60
taatggaatc	ctctagggtt	aagccaaaga	aaaattggag	agacaaaatt	agattttgta	120
gcccttttcc	cccccgga	tgcccttttt	tttcttttta	gtttctgatg	aatggctatc	180
atztatttct	accaaattta	aataaggact	gctgccttgt	atgtttaact	aggcaggcag	240
agggaactgg	tttgtttagg	aagcagtgac	tgagatgtcc	tgccaagtt	agtgcacagag	300
gaggggagaa	agaatccaga	ccaatttgta	tgagtatat	tttactcca	tgaaataaaa	360
cacatttggt	tcatatttgc	tgaaaagtaa	aacaataata	ttgtacgaaa	tgttatacac	420
agggtagggt	gtacatagca	gtttcagaaa	catcattgca	tccaccagag	aaactattct	480
aaaactgata	ttcacacatt	ttttataata	ataataatat	gttagaaaca	tacagtgtgg	540
catttagtat	atacactccc	ttgctcgcaa	gcgaaaaatc	ctaactcgct	ctgtataaca	600
tgctttat	taaagcctaa	cctttaaaaa	cactgttgtg	atattactaa	caactgcttt	660
tataaaatta	atttgacatt	tcgatataata	tacatccttt	cagtcattta	aatgttaaca	720
atgctaaact	taaaaaataa	caagcttata	gtaatgttaa	aatgtcatat	ccagtcacaa	780
atgtgtgtgt	gtatgtgtcc	ttgcaactgt	tagaaatact	tgtagtgaaa	gatgtcagac	840
actgaggaca	tccctttgaa	atcaaaggag	ctctctcttt	gattcagtg	tttccttttc	900
tctatatagc	ttctctttct	ctccctttct	ttagtgccca	cgaccttcta	gcataattcc	960
cagtctttca	agggcgaggt	tgccccatcc	ggcaagggtc	taggatcccg	gcgctgtggg	1020
tgcggtctac	acgggcccgt	ccactgcata	ctggcaagca	ctcaggttgg	aggccgggtt	1080
ctgcacgctg	gcgtagccga	agctggagtg	ctgctttgct	ttcagttcca	ggctggccag	1140
gctcgagtta	cacgtgtccc	tataaacata	cggaggagtc	ggcggcgcgt	aaggacaggc	1200
aggcgctcgg	acgcggaat	tcagcgacgg	gctactcagg	ttgttcaagt	tattcaggct	1260
gttgagactg	gagcccggga	cgcctgtcac	tgctgagggc	accatgctgg	acgacatgct	1320
catggacgag	atagagttgg	gtggggaaaa	catgctctgt	gatgacaggg	gggtgacgtt	1380
catagagttg	aagaagggga	agctcttggt	ggatagggag	gcggatgtaa	ggcccttggc	1440
ggcccagttg	ttgtaggaat	agcctgggta	catgtcgtcg	tagggctgca	tgagcccatt	1500
gaactgcggc	ccgaagccat	tcttgcatag	ctcgccctgc	tggttgcgct	ccctctttct	1560
ccatttggcc	cgacgattct	tgaaccaaac	ctgggggagg	ttggggcaag	ggagcaaaaca	1620
gatgccacag	tgcagattac	taaaacttcc	atcggaggcc	aacccccgcc	ttcccccgac	1680
acacacgcta	gcgcactcac	acaccctggc	ctcgcttcac	tgacccgccc	tgcaacacaa	1740
gataccaggg	ccagctttca	gttactggcc	cgggtctcca	ccaagcgcag	gagacctggt	1800
ctgctctggc	ctgcgagctg	ggactcggag	ctacgccaca	aacctcagcc	gaacgcattg	1860
agacctcg	acggtttgat	cactcagcca	ggcgtttctc	caggtccaaa	aacacttaat	1920
gtaaaacaaa	cgcggggag	caggcttttc	caacccttcc	cggggcacct	tgcaaaactg	1980
cttccattcc	aaagccacag	acccacggat	gaggagaagg	ggctggaagg	gcactagagg	2040
atcgctcttt	ctcccacgca	attcctccct	tccttccctg	acctccactg	tcgtccccc	2100
ccccctggta	cgtgctccct	taacaggggac	taggcgcgca	acactctttc	tcgcctagca	2160
aaacaaccaa	ataaagagca	aaagaccacc	tcttgcgtcag	ctcgtttaact	ccaggagctt	2220
ggcatattaa	actccgggaa	cccggaagg	gtagtttttg	agattccccc	ttctttcgct	2280
ctgcctcttc	tttaccctaa	gccaccaca	ggcctgtccg	cgcgccaggc	ccagccgggt	2340
cgtttggtt	tgcaggcgcc	caccagggc	ggccggcttc	cacccgtgtc	cggtggccca	2400
gccgcaaccc	cgatcccaat	ccacatcg	cctccctgtc	gccccagacg	gcggcttttg	2460
tgtattggag	agaggcctgg	cctgagatat	ccgagctgac	accagtgatg	tttcacatta	2520
cacatctccg	ccggggccag	ccgtgtaatc	cgctttttct	ctttttcctt	tcattcttga	2580
tttccttttt	atcccccttc	ctctttgcac	ccgactgcta	taaaaagcac	gcctcactcc	2640
cacttggtc	gacaagcagc	cgccctggaa	ggagaggcag	ctgcaaggag	agcccagcgc	2700
cgcggctaca	aagcactagg	gtggagctgc	ggaatagcgg	gcgggggtgg	agggcgtttt	2760
cgaaggatcc	cagaaaaccc	atagactctg	tctttaatta	cttgccattt	ctaccctagg	2820

ccatctaaac	tttgctcagg	cgagaagagt	acgtgagagg	cccgttccct	tgatgtgcaa	2880
gagagcta	gaaagactga	ccttgctcaa	aaccacgccg	cccaggaccc	agctctggct	2940
ctggacagtt	aaactaaaac	cattttcaac	ttcttcccgg	ccttttatcc	accagcatag	3000
cctcatgcct	tgacacaaatg	ccaccagag	agtgtcttca	ttccctctga	tttgggagag	3060
cattttggtc	tttattcttt	ttatcgttgt	tttcttcttt	ttgtttgctc	tgctctaacc	3120
gggggcttta	ttttttctac	ccagagcact	taattttttt	tttttaacag	caaagcctct	3180
ggatgccgct	tgatttgctt	gattctgttt	tctgcttcca	gaatcctaac	aaatttgga	3240
tcttccaccg	accagcataa	accaggacgt	tgctattggg	ttattttatt	gagctcattt	3300
ttgccaatcc	ataaagtaca	gatttgctac	aaagttaagg	taagcccttt	ttacaaaact	3360
atgattataa	tttagaagag	ggggtgtgag	tttcaatttc	cagagttcaa	ctcctgagag	3420
aagataaata	aaccaagcag	aaaagtcttt	cttctttttt	tctttctcct	tctaagagga	3480
ctagtagttg	tgtattaaaa	ctttgctccc	ggagatcaca	aaactaggaa	ataggggtgtg	3540
tgggagagac	ctgaatggcc	gaaacaaccg	taaagaaggt	gtaagaagcg	cgagcccagg	3600
agggaaaaag	ctggggccagg	gccgggacaa	aggtttccca	gggagggccca	actcttccgt	3660
gtctctggcg	ggttttccct	gttaaaggct	cacaggttgg	agcctgttcg	cggctcttgg	3720
cctggtaggg	attttattag	ctctgctctg	gcaactgcaa	gccaggaaaca	caatgtcctg	3780
tgcaggggat	tgcccatgca	gcccagctcg	tgagatcgcg	ggatggcggg	gcagtgagcc	3840
ggtgccgctc	tgggagcctg	agccaggggcg	cgagtcctgt	cggcctcgga	gagggaaactg	3900
taatctcgca	accaggccgc	cgcgaggcct	tctgcctttg	caaagctgcg	ccccaccggc	3960
gccctcccag	gcggcgctgc	cttccacatt	ctctcctggg	ctacttgccc	tgtacctcca	4020
caacatcctc	cccccatccc	tcccagactc	cgtgctggct	cctaccggga	ctcgggcttc	4080
cgtaaggttg	gtccacacag	cgatttcttc	gcgtgtggac	atgtccgggt	agcgggttcc	4140
ctggaaagtg	gcctccagct	cctggagctg	ctgggtggta	aagttagtcc	gctgccgcct	4200
ttgccgcttc	ttcttagacg	ggctcctggc	gcccacgtcc	tcattcttcc	cctgctggct	4260
tttatctttc	tctgaaaacg	aaacacacac	actttcccgt	cagcatgccc	acctgcaacg	4320
cggacgcaa	ctggaccggc	ggcagaagcc	gtggaaagag	tgggctgcct	ggcgccggag	4380
gaggggtgcg	cggcgcgctc	cgggcccgcga	ggagcgctgc	gcctgtgggg	tgtgcaggcg	4440
caagtgtggg	tgtccgcgcc	ccatttcttc	ccctccccca	gcgcccacg	ttttattttac	4500
atgtttatct	cactgcagcg	gcacattcac	ttttatagcc	tgtgctttca	agtatatatta	4560
tacacctctg	cgacagacac	ccaaatctcc	tgggacgcgc	acacgcgcgt	ggtttacaga	4620
ccccctccc	cctcgcagaa	agctcagatt	tccatgcggt	ttgggaaggc	taggaaaaga	4680
tgtggggatt	cgggttgggca	ccgaagttcg	ccggcccttt	cccaaaaaaa	aaaaaaaaat	4740
gcctcttcgc	gaagggcatt	tctgagtggg	ttcaggcaat	ttcctaacga	gtggagctcc	4800
tcgggagctg	aaagccgaga	ggaaaacagg	gacagaggtc	ggcgccctct	gaagctcctc	4860
gaatcaagat	gctgggattt	ttgtgaccca	ggaaaacagaa	gggaggccag	ggtacgaata	4920
gagaggcgcg	cagaattgct	cgcgccctta	gcgccccagg	agccggggcg	gtcgaggggag	4980
aactaaaggg	atgcggggta	gtcaaaaattc	cggctcccgg	aagttctgcg	gggagccagg	5040
cgaacgacca	ctcccaccac	gcctcccccc	ggaggggctg	acttccttgg	ggcgagaggg	5100
agcgggtggc	gcagagcagc	tgagcgggaa	tgtctgcagg	gcggcgcggc	gccttacctg	5160
cggcctccgg	gctggaggtg	tcggagatgg	tgtgcacctc	cagcctgtgc	ttggaggagt	5220
ccagcgaccg	gggtgaccg	ggagccagaa	ccgaagccat	ggctaacggc	tgggatggg	5280
gacaggaaga	tgaggagacg	gccgacagct	tggtccccgc	tgctcgggtg	tccaagtga	5340
gcgggccttt	catgcagttc	atggacgag	gagcgcgacg	ctctactagt	ccttggtac	5400
tgcccccgct	agcccccgta	gccgcgctg	cccgctccgg	gtcgcgctct	aggcgcgag	5460
tttccccgct	gcggggagag	ccaggggacg	caacccccgc	cgagttctca	agccaagctg	5520
cccccgctct	ctccggaagg	ctcaagcgaa	aaagtccgga	gacggaaagt	cagcgggcaa	5580
acgaagacat	gggatgtggg	cagaagggca	ccactcagag	cgtcttttag	gagcaggctt	5640
ccaagctcca	aagcgaaaca	agagtgggca	aagaccccc	tcttctctcc	ctccctcccc	5700
caagaacccc	tccaataagg	aaagctaacg	ccgaccgcgc	tctgcccggc	ccccccccac	5760
gcggcagccc	tgacagagaa	gtgtcaagag	tgacagggac	aggtaggtga	tattagatcc	5820
cctgcggcgg	cagcagccgc	tgcagccacg	acgcggccct	ctgagcgcac	cctccgcaac	5880
gcgcacacgc	acacccctcg	ggcgggtcgaa	caggagccgg	gccttgccgc	agctcagctc	5940
caggcaccca	ggcgagcgac	ggaccagatc	tgcggctccg	cgcttccctg	ttggcctaac	6000
atcttaaaac	cagaggcggg	cttctgtgtg	ccgagacgtc	actccgccgc	ggccctcccc	6060
agccctctcc	gcctccgcct	cctcccagac	ccttctccgg	gtgcgactga	cgtgggtccg	6120
caccaatcag	gacgccccga	gccgcgggtg	agggactgtc	ctgcctgcac	ctatcagcag	6180
tgcggggccg	ggctactgoc	tcgcogtgcg	cactgggtct	acacaggcaa	gctcccggga	6240
attcagctcc	tgcccagccc	aaggcgatcc	ggcttttagt	acgaacccaa	aggtgaagag	6300
atgaggctag	gagtcgaagg	cttggggagaa	gagagtggaa	tggtaagaa	gagaaaagga	6360
caaggatcaa	caagacaccc	actcttttgt	tctcactaca	tccattttcca	atccccacc	6420
ccatataaaa	aggagacacg	ttacttaaaa	ctagaaaatt	tgaaaaacag	caacaaatca	6480
cctctccgat	cttaaatctt	ccaaacagcc	tgtcaagtga	atgctgcgct	aatctgaaga	6540
agctttaatt	gcaaagaaga	cagagccctg	aaaaggcagg	ctaataaatt	agaaatcgag	6600

aagcaaattgg	acccgtcaaa	agaaaattac	cttgacttta	aacgaacaac	tgtttggtgg	6660
ttcactctgg	atttatacaa	gaataaaaag	tgcctcaga	tcacgttctc	tgtgatgctt	6720
attagtcccc	agacagaaaa	cacacaatag	aagagaaacc	ctaaccagc	gttttcaaaa	6780
tgctgaaagc	ttatccattc	tacttaacgt	tgattaagac	acatatccta	gatctttcaa	6840
attccttgta	cactgtatta	agctcgtcct	aacccgagag	agccacgctt	taaattcgac	6900
tctcttgttt	actttattat	caatcagatt	taaatccata	aagcctgtag	aatcaacaac	6960
cttgagctaa	ttatatatga	aatatgcctt	aatgaatttc	catacaatta	agaatgttgc	7020
caaataacca	atttcaagga	taatttttaa	cagtcatttt	cttttcccag	tgagctcaag	7080
gctgtcttga	gccattaaag	tccaagcagg	cagaaggggt	gtgtgtgagc	taaggggcga	7140
aagcctagaa	ctgcgctcaa	ctagcaaaaag	caaaacctta	tttatataaa	acaaaaaa	7200
tcacctttgg	agacatcaac	tctttatagc	actgtttcca	agcaaattta	attttccaag	7260
aaattaaaga	aagaaatcca	aacatattca	aaataatttt	tgaaagtcct	tttgtcccc	7320
agcataggtc	agctggagag	gacaaaactaa	tctcctctgg	gtttctgcat	gggcgattgt	7380
tttactatgg	agttagtgtt	atcatctctg	aatgtgtatt	tgtttgacat	tacagtcaat	7440
gatttgcaat	gccagcatga	agtatcttta	aaacactccc	tccttgctct	tgttcacaag	7500
attgggaaac	ttattcgaatg	tggaacaaag	tggatgaagc	agactacaaa	tatatattgca	7560
acttatgtgt	ctctcctttg	ccctgaccac	ccccaaaacc	tatctgcaac	tcctcccat	7620
tttaaacttg	cagtcctaaag	acgcacatga	gaattgtttt	tcagtctttc	ttcaccagta	7680
tcaccccact	ttagaataaa	tttagctgca	agggaggaat	ttcttcatag	taagcttttaa	7740
atcagcattt	ctgcttttaa	ccttttatct	cactttaccc	cattccacac	atcacagacac	7800
ctgctcagag	taaaacacat	cctcatgtga	caggtctgca	ttagctgagg	ctcatagac	7860
cagctatatt	aggtcctgca	atcttatcac	taaattatac	acattacact	agcagcctgt	7920
tggtaaagaa	ggttaaatta	attttacattc	tgctcattat	ctggtgctta	aatgaogcat	7980
tttatcccgg	agatttggtg	gagaatctcc	ttctcagacc	ccacagcgtt	tcactgaaga	8040
caatgctctt	acatttgtag	tggttttttaa	tctgataaga	ctctaatttg	cttaagtctt	8100
ttaaataaag	gttttaaatg	tttctagccg	ttttcttatt	gaatttcctc	taattccccc	8160
aagatcataa	agtatatgtg	taaagtaaat	atttccctcc	attgcaactgc	cagccgatga	8220
cctataacta	agtcaataag	aatccagctc	ttttctgctg	aatgtgttta	ctaatacatat	8280
tccagtttct	tcttttaaac	ctcagaatag	ctgtggtccc	cacaatacca	tgccccttaa	8340
agcttcattt	catgaaggga	ctccatcaca	ttaaagaatg	aaaaaaatct	ccactgtagt	8400
tagtatacac	agcccctcac	tccttggttt	tcaagattca	aacccagag	ctgcaaatat	8460
ttttggaagc	ttgggtgtta	atgccttatt	ttagaaagcc	gagaagcccc	acagagccat	8520
atagatttct	aaacccatct	ctcataaaac	cacagaattt	tgataaaaagc	tctggtggct	8580
ccactctacc	gatggaactt	tcacacagac	aaatatacat	gtatgaagga	cctcaatcag	8640
cctccaaagt	ggttgaaaaa	cccaagggca	cgtgactgct	cctcatagt	ccaacgtgtg	8700
cgagatgttg	gaagcactgg	ggatcagcag	cagcctagat	gcctaaaaag	ataaggtgtc	8760
ctaatttggt	tggaccattt	gaagtcaagt	gggtgaataaa	gacaattatc	tagataattc	8820
agattaaagt	aaaagcaaaa	ccatatctat	ttgtatatat	atattcacat	ccattttgata	8880
ttacagacac	acacacgcac	acacacactg	gctctgtaaa	caactgactc	aaagttagga	8940
ttttctttgc	atttttccag	caggagtttc	aacattctcc	taatctccta	atcactttac	9000
a						9001

<210> 2

<211> 9001

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 2

agttgatgga	tttggttaa	ttttttttt	ttttttttt	tttatattat	ttgttagtta	60
taatggaatt	tttttaggtt	aagttaaaga	aaaattggag	agataaaatt	agattttgta	120
gtttttttt	ttttcgggaa	tggtttttt	ttttttttt	gtttttgatg	aatggttatt	180
atttttttt	attaaattta	aataaggatt	gttggtttgt	atgttttaatt	aggtaggtag	240
agggaattgg	tttggttagg	aagtagtgat	tgagatgttt	tggttaagtt	agtgatagag	300
gaggggagaa	agaatttaga	ttaatttgta	tgtagtatat	ttttttttt	tgaaataaaa	360
tatatattgt	ttatatattg	tgaaaagtaa	aataataata	ttgtacgaaa	tggtatatat	420
agggtagggt	gtatatagta	gttttagaaa	tattattgta	tttattagag	aaattatttt	480
aaaattgata	tttatatatt	ttttataata	ataataatat	gttagaaata	tatagtgtgg	540
tatttagtat	atatattttt	ttgttcgtaa	gcgaaaaatt	ttaatcgttt	ttgtataata	600
tgttttattt	taaagttaa	tttttaaaaa	tattgttgtg	atattattaa	taattgtttt	660

tataaaatta	atttgatatt	tcgatatata	tatatTTTTT	tagttatttta	aatgttaata	720
atgttaaatt	taaaaaataa	taagtttata	gtaatgttaa	aatgttatat	ttagttaaat	780
atttgtttgt	gtatgtgttt	ttgtaattgt	tagaaatatt	tgtagtgaat	gatgttagat	840
attgaggata	tttttttgaa	attaaaggag	tttttttttt	gattttagtgg	tttttttttt	900
tttatatagt	tttttttttt	tttttttttt	ttagtgttta	cgatttttta	gtataatttt	960
tagtttttta	agggcgagg	tgtttttatt	ggtaagggtt	taggatttcg	gcgttgtggg	1020
tgcggtttat	acgggtcggt	ttattgtata	ttggtaagta	tttaggttgg	agggtcggtt	1080
ttgtacgttg	gcgtagtcga	agttggagtg	ttgttttgtt	tttagtttta	ggttgggttag	1140
gttcgagtta	tacgtgtttt	tataaatata	cggaggagtc	ggcggcgctg	aaggataggt	1200
aggcgctcgt	atcgcggaat	ttagcgacgg	gttatttagg	ttgtttaagt	tatttagggt	1260
ggtgagattg	gagttcgga	cgtttggtat	tggtgagggg	attatgttgg	acgatatggt	1320
tatggacgag	atagagttgg	gtggggaaaa	tatgttttgt	gatgataggg	ggttgacggt	1380
tatagagttg	aagaagggga	agtttttggt	ggatagggag	gcggatgtaa	ggtttttggc	1440
ggttttagttg	ttgtaggaat	agtttgggta	tatgtcgtcg	tagggttgta	tgagttttat	1500
gaattgcggt	tcgaagttaa	ttttgtatag	ttcggtttgt	tggttgcgtt	tttttttttt	1560
ttatttggtt	cgacgatttt	tgaattaaat	ttggggcggt	ttggggtaag	ggagtaaata	1620
gatgttatag	tgtagattat	taaaattttt	atcgagggtt	aattttcggt	ttttttcgat	1680
atatacgtta	gcgtatttat	atatatttgt	ttcgttttat	tgtatcggtt	tgtatattaa	1740
gatattaggg	ttagttttta	gttattgggt	cgggttttta	tttagcgtag	gagatttggt	1800
ttgttttggt	ttgcgagttg	ggattcggtg	ttacgttata	aattttagtc	gaacgtatgg	1860
agatttgctg	acgggttgat	tatttagtta	ggcgtttttt	taggttttaa	aatatttaat	1920
gtaaaataaa	cgcggggtag	taggtttttt	taattttttt	cgggggtatt	tgtaaatttg	1980
tttttttttt	aaagttatag	atttacggat	gaggagaagg	ggttggaagg	gtattagagg	2040
atcgtttttt	tttttacgta	attttttttt	tttttttttt	attttttatt	tcgtttttta	2100
tttttttggt	cgtgtttttt	taatagggat	taggtcgtta	atattttttt	tcgttttagta	2160
aaataattaa	ataaagagta	aaagatttat	ttttcgttag	ttcgttaatt	ttaggagttt	2220
ggatatattaa	atttcgggaa	ttcggaaagg	gtagttttgg	agattttttt	ttttttcgtt	2280
ttgttttttt	tttattttta	gtttattata	ggtttgttcg	cgcgtttagg	ttagtcgggt	2340
cgtttggttt	tgtaggcggt	tatttaggtc	ggtcgggttt	tattcgtgtt	cgggtggttt	2400
gtcgttaatt	cgattttta	ttatatcggt	tttttttgct	gttttagacg	gcggtttttg	2460
tgtattggag	agaggttttg	tttgagatat	tcgagttgat	attagtgatg	ttttatatta	2520
tatattttcg	tcgggttttag	tcgtgtaatt	cgtttttttt	tttttttttt	ttatttttga	2580
tttttttttt	attttttttt	tttttttgat	tcgattgtta	taaaaggtac	gtttttattt	2640
tatttggttc	gataagtagt	cgtttttgga	ggagaggtag	ttgtaaggag	agtttagcgt	2700
cgcggttata	aagtattagg	gtggagttgc	ggaatagcgg	gcgggggtgg	agggcggttt	2760
cgaaggattt	tagaaaattt	atagattttg	tttttaatta	tttggtattt	ttatttttag	2820
ttatttaaat	tttggttttag	cgagaagagt	acgtgagagg	ttcgtttttt	tgatgtgtaa	2880
gagagttaat	gaaagattga	ttttgtttta	aattacgtcg	tttaggattt	agttttggtt	2940
ttggatagtt	aaattaaaa	tattttta	tttttttcgg	ttttttattt	attagtatag	3000
ttttatgttt	tgtataaatg	ttatttagag	agtgttttta	ttttttttga	tttgggagag	3060
tatttttggt	tttatttttt	ttatcgttgt	tttttttttt	ttgtttgttt	tgttttta	3120
gggggtttta	ttttttttat	ttagagtatt	taattttttt	tttttaatat	taaagttttt	3180
ggatgtcgtt	tgatttggtt	gattttgttt	tttggtttta	gaattttaat	aaatttgga	3240
ttttttatcg	attagtataa	attaggacgt	tgttattggg	ttattttatt	gagttttatt	3300
ttgttaattt	ataaagtata	gatttgttat	aaagttaagg	taagtttttt	ttataaaatt	3360
atgattataa	tttagaagag	ggggtgtgag	tttttaattt	tagagttaaa	tttttgagag	3420
aagataaata	aattaagtag	aaaagttttt	tttttttttt	tttttttttt	tttaagagga	3480
ttagtagttg	tgtattaaaa	ttttgttttc	ggagattata	aaattaggaa	atagggtgtg	3540
tgggagagat	ttgaatggct	gaaataatcg	taaagaaggt	gtaagaagcg	cgagtttagg	3600
agggaaaaag	ttgggttagg	gtcgggataa	aggtttttta	gggagggtta	attttttcgt	3660
gttttttggt	ggtttttttt	gttaaagggt	tatagggttg	agtttgttcg	cgggtttttg	3720
tttggtaggg	attttattag	ttttgttttg	gtaattgtaa	gttaggaata	taatgttttg	3780
tgtaggggat	tgtttatgta	gttttagttc	tgagatcgcg	ggatggcggt	gtagttagtc	3840
ggtgtcgttt	tgggagtttg	agtttagggc	gtagttttgt	cgggttcgga	gaggggaattg	3900
taatttcgta	attaggtcgt	cgcgagggtt	tttggttttg	taaagtgtcg	ttttatcggc	3960
gttttttttag	gcggcggttg	tttttatatt	tttttttggt	ttatttggtt	tgtattttta	4020
taatatattt	ttttattttt	tttttagatt	cgtgttggtt	tttttcgga	ttcgggtttt	4080
cgtaagggtg	gtttatatag	cgattttttt	gcgtgtggat	atgttcgggt	agcgggtttt	4140
ttggaaagtg	gttttttagt	tttgaggttg	ttgggttgga	aagttagttc	gttgcgtttt	4200
ttgtcgtttt	tttttagacg	ggtttttcgg	gtttacgttt	ttattttttt	ttgttggtt	4260
tttatttttt	tttgaaaacg	aaatatatat	attttttcgt	tagtatgttt	atttgtaacg	4320
cggacgttaa	ttggatcggc	ggtagaagtc	gtggaagagt	tgggttggtt	ggcgtcggtg	4380
gaggggtcgc	gcggcggttt	cgggtcgcga	ggagcgttgc	gtttgtgggg	tgtgtaggcg	4440

taagtgtggg	tggtcgcgtt	ttattttttt	ttttttttta	gcgtcgtacg	ttttattttat	4500
atgtttat	tattgtagcg	gtatat	ttttatagtt	tgtgtttttt	agtatat	4560
tatat	cgtagatata	ttaaat	tgggacgcgt	atacgcgcgt	ggtttataga	4620
ttttttttt	tttcgtagaa	agtttagatt	tttatgcggt	ttgggaaggt	taggaaaaga	4680
tgtggggatt	cgggtgggta	tcgaagttcg	tcggtttttt	tttaaaaaaa	aaaaaaaaat	4740
gttttttcgc	gaagggtatt	tttgagtgg	tttaggta	tttttaacga	gtggagtttt	4800
tcgggagttg	aaagtgcgaga	ggaaaatagg	gatagaggtc	ggcgggtttt	gaaggttttc	4860
gaattaagat	gttgggattt	ttgtgattta	ggaaatagaa	gggaggttag	ggtacgaata	4920
gagagggcgg	tagaattggt	cgcgttttta	gcgttttagg	agtcgggtcg	gtcaggggag	4980
aattaaaggg	atgcggggta	gttaaaat	cggttttcgg	aagttttg	gggagttagg	5040
cgaacgatta	tttttattac	gttttttttc	ggaggggttg	atttttttgg	ggcgagaggg	5100
agcgggtggc	gtagagtagt	tgagcgggaa	tggttgtagg	gcggcgcggc	gttttatttg	5160
cggttttcgg	gttggaggtg	tcggagatgg	tgtgtatttt	tagtttggtg	ttggagaggt	5220
ttagcgatcg	gggttgatcg	ggagttagaa	tcgaagttat	ggttaacggt	tggggatgg	5280
gataggaaga	tgaggagacg	gtcgatagtt	tggttttcgt	tgttcgggtg	tttaagtga	5340
gcgggttttt	tatgtagttt	atggacgagg	gagcgcgacg	ttttattagt	ttttggttat	5400
tgtttcgtcg	agttttcgt	gtcgtcgttg	ttcgtttcgg	gtcgcgtttt	aggcgcggag	5460
ttttttcgtt	gcggggagag	ttaggggacg	taattttcgt	cgagttttta	agtttaagttg	5520
ttttcgtttt	tttcggaagg	tttaagcgaa	aaagttcgg	gacggaaagt	tagcgggtaa	5580
acgaagatat	gggatgtggg	tagaagggt	ttatttagag	cggttttagg	gagtaggttt	5640
ttaagtttta	aagcgaaata	agagtgggt	aagatttttt	tttttttttt	tttttttttt	5700
taagaatttt	tttaataagg	aaagttaacg	tcgatcgcgt	tttggttcgt	ttttttttac	5760
gcggtagttt	tgatagagaa	gtgttaagag	tgatagggat	aggtaggtga	tattagattt	5820
tttgcgcgcg	tagtagtcgt	tgtagttagc	acgcgggttt	ttgagcggt	ttttcgtaac	5880
gcgtatacgt	atatttttcg	ggcgggtcgaa	taggagtcgg	gttttgctgt	agtttagttt	5940
taggtattta	ggcgagcgac	ggattagatt	tgcgggttcg	cgtttttttg	ttgggttaaat	6000
attttaaaat	tagaggcggg	ttttttggtg	tcgagacggt	atttcgtcgc	ggtttttttt	6060
agtttttttc	gttttcgttt	tttttttagt	tttttttcgg	gtgcgattga	cgtgggttcg	6120
tattaattag	gacgtttcga	gtcgcgggtg	agggattggt	ttgtttggt	ttattagtag	6180
tgcggggtcg	ggttattggt	tcgtcgtg	tattgggttt	atataggtaa	gttttcggga	6240
atntagtttt	tgtttagttt	aaggcgattc	ggtttttagt	acgaatttaa	aggtgaagag	6300
atgaggttag	gagtcgaagg	tttgggagaa	gagagtggaa	tggttaagaa	gagaaaggta	6360
taaggattaa	taagatat	attttttg	ttttattata	tttattttta	attttttatt	6420
ttatataaaa	aggagatacg	ttatttaaaa	ttagaaaatt	tgaaaaatag	taataaatta	6480
ttttttcgat	tttaaat	ttaaatagtt	tgtaagtga	atgttgcggt	aatttgaaga	6540
agttttaaat	gtaaagaaga	tagagttttg	aaaaggtagg	ttataaaat	agaaatcgag	6600
aagtaaatgg	attcgttaaa	agaaaattat	tttgatttta	aacgaataat	tgtttggtgg	6660
tttatttttg	atttatataa	gaataaaaag	tcgttttaga	ttacgttttt	tgtgatgttt	6720
attagttttt	agatagaaaa	tatatataat	aagagaaatt	ttaatattagc	gttttttaaaa	6780
tgttgaaagt	ttattttatt	tatttaacgt	tgattaaagt	atatattttta	gatttttttaa	6840
attttttgta	tattgtatta	agttcgtttt	aatttcgagag	agttacgttt	taaattcgat	6900
ttttttgttt	attttattat	taattagatt	taaatttata	aagtttgtag	aattaataat	6960
tttgagttaa	ttatatatga	aatatgtttt	aatgaatttt	tatatataat	agaatgttgt	7020
taaataatta	attttaagga	taatttttaa	tagttatttt	tttttttttag	tgagtttaag	7080
gttgttttga	gttattaaag	tttaagttag	tagaaggggt	gtgtgtgagt	taagggcgaa	7140
aagtttagaa	ttgcgttttaa	ttagtataag	taaaatttta	tttatataaa	ataaaaaaaa	7200
ttatttttg	agatat	tttttatagt	attgttttta	agtaaaattta	attttttaaag	7260
aaattaaaga	aagaaattta	aatatat	aaataatttt	tgaaagtttt	ttgttttttt	7320
agtataggtt	agttggagag	gataaa	tttttttttg	gtttttgtat	gggcgattgt	7380
tttattatgg	agttagtgtt	attatttttg	aatgtgtatt	tgtttgatat	tatagttaat	7440
gatttgta	gttagtatga	agtattttta	aaatat	tttttg	tgtttataag	7500
attgggaaat	ttattcgatg	tggaataaag	tggatgaagt	agattataaa	tatatattgta	7560
atttatgtgt	tttttttttg	ttttgattat	ttttaaat	tatttgta	ttttttttat	7620
tttaaaattg	tagtttaag	acgtatatga	gaattgtttt	ttagtttttt	tttattagta	7680
ttatttttatt	ttagaataaa	tttagttgta	agggaggaat	ttttttatag	taagtttttaa	7740
attagttatt	ttgttttttaa	ttttttattt	tattttatat	tattttatat	atatagatat	7800
ttgttttagag	taaaatata	ttttatgtga	taggtttgta	ttagttagg	ttatatatt	7860
tagttatatt	aggttttgta	attttattat	taaattatat	atattatatt	agtagtttgt	7920
tggttaaagaa	ggttaaatta	attttatatt	tgtttattat	ttggtgttta	aatgacgtat	7980
tttattttcgg	agatttg	gagaattttt	tttttagatt	ttatagcgtt	ttattgaaga	8040
taatgttttt	atatttgtag	tgggtttttaa	tttgataaga	ttttaatttg	tttaagtttt	8100
ttaaataagg	gttttaaatg	tttttagtcg	tttttttatt	gaattttttt	taattttttt	8160
aagattataa	agtatatgtg	taaagtaaat	attttttttt	attgtattgt	tagtcgatga	8220

tttataatta	agttaataag	aatttagttt	ttttttgttg	aatgtgttta	ttaattatat	8280
tttagttttt	ttttttaaat	tttagaatag	ttgtggtttt	tataatatta	tgttttttta	8340
agttttat	tatgaaggga	ttttattata	ttaaagaatg	aaaaaaat	ttattgtagt	8400
tagtatatat	agttttttat	tttttgtttt	ttaaagattta	aatttttagag	ttgtaaatat	8460
ttttggaagt	ttgggtgtta	atgtttttat	ttagaaagtc	gagaagtttt	atagagttat	8520
atagattttt	aaatttat	tttataaatt	tatagaattt	tgataaaaagt	tttgggtggt	8580
ttattttatc	gatggaattt	ttattacgat	aaatatatat	gtatgaagga	ttttaattag	8640
tttttaaagt	ggttgaaaaa	tttaagggtta	cgtgattggt	ttttatagt	tttaacgtgtg	8700
cgagatgttg	gaagtattgg	ggattagtag	tagtttagat	gtttaaaaag	ataaggtggt	8760
ttaatgtgtg	tggattttat	gaagttaagt	ggtgaataaa	gataattatt	tagataattt	8820
agattaaagt	aaaagtaaaa	ttatattttat	ttgtatatat	atattttatat	ttattttata	8880
ttatagatat	atatacgtat	atataatttg	gttttgtaaa	taattgattt	aaagtgagga	8940
ttttttttgt	atttttttag	taggagtttt	aatttttttt	taatttttta	attattttat	9000
a						9001

<210> 3

<211> 9001

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 3

tgtaaagtga	ttaggagatt	aggagaatgt	tgaaattttt	gttggaaaaa	tgtaaagaaa	60
attttttatt	tgagtttagt	gtttatagag	ttagtgtgtg	tgtgcgtgtg	tgtgtttgta	120
atataaaatg	gatgtgaata	tatatatata	aatagatatg	gttttgtttt	tattttaatt	180
tgaattat	agataattgt	ttttatttat	tatttgattt	taatgggttt	atataaatta	240
ggatatttta	tttttttagg	tatttagggt	gttggtgatt	tttagtggtt	ttaatatttc	300
gtatacgttg	gtattatgag	gagtagttac	gtgttttttg	gttttttaat	tattttggag	360
gttgattgag	gttttttata	tatgtatatt	tgtcgtgatg	aaagttttat	cggtagagtg	420
gagttattag	agttttttat	aaaattttgt	gggtttatga	gagatgggtt	tagaaaattt	480
tatggttttg	tgggggtttt	cgttttttta	aaaaataagg	ttaatattta	agtttttaaa	540
aatatttgta	gttttggggg	ttgaattttg	aaaaataagg	agtgggggtt	tgtgtatatt	600
aattatagt	gagatttttt	ttatttttta	atgtgatgga	gtttttttat	gaaatgaagt	660
tttaaggggt	atggtattgt	ggggattata	gttattttga	ggtttaaaag	aagaaattgg	720
aatatgatta	gtaaatatat	ttagtagaaa	agagttggat	ttttattgat	ttagttatag	780
gttatcgggt	ggtagtgtaa	tgggaggaaa	tatttatatt	atatatatat	tttatgattt	840
tgggggaatt	agaggaaatt	taataagaaa	acggttagaa	atattttaaa	tttttattta	900
aaagatttaa	gtaaattaga	gttttattag	attaaaaatt	attataaatg	taagagtatt	960
gttttttagtg	aaacgtttgt	gggttttgaga	aggagatttt	tcgttaaaat	tctgggataa	1020
aatgcgttat	ttaagtatta	gataatgagt	agaatgtaaa	ttaatttaat	tttttttatt	1080
aataggttgt	tagtgtaatt	tgtataattt	agtataaga	ttgtaggatt	taatatagtt	1140
ggatgttatga	gttttagtta	atgtagattt	gttatatgag	gatgtgtttt	attttgagta	1200
ggtgttttga	tgtgtggaat	ggggtaaaagt	ggaataaaaag	gttaaaagta	gaaatgttga	1260
tttaaagttt	attatgaaga	aatttttttt	ttgtagttaa	attattttta	aagtgggatg	1320
atattggtga	agaaagattg	aaaaataaatt	tttatgtgcg	tttttggatt	gtaagtttaa	1380
aatggggagg	agttgttagat	aggggtttggg	ggtggttagg	gtaaaggaga	gatataataag	1440
ttgtaaatat	atttgtagtt	tgtttttatt	attttgtttt	atatcgaata	agttttttta	1500
ttttgtgaat	aaggataagg	agggagtgtt	ttaaagatat	tttatgttgg	tattgtaaat	1560
tattgattgt	aatgttaaat	aaatatatat	ttagagatga	taatatataat	tttatagtaa	1620
aataatcgtt	tatgtagaaa	tttagaggag	attagtttgt	tttttttagt	tgattttatgt	1680
tgggggataa	aaggattttt	aaaaattatt	ttgaatatgt	ttggattttt	ttttttaatt	1740
tttttggaaa	ttaaatttgt	ttggaaatag	tgttataaag	agttgatgtt	tttaaagggtg	1800
attttttttg	ttttatataa	ataagggttt	gtttttgtta	gttgagcgta	gttttagggt	1860
tttcgttttt	agtttatata	tatttttttt	gtttgttttg	attttaaatg	tttaagatag	1920
ttttgagttt	attgggaaaa	gaaaatgatt	gttaaaaaatt	atttttgaaa	ttggttattt	1980
ggtaaatatt	ttaatgtgat	ggaaatttat	taagggtata	tttatatat	attagtttaa	2040
ggttgttgat	tttatagggt	ttatggattt	aaatttgatt	gataataaag	taaaataagag	2100
agtccaattt	aaagcgtggt	tttttcgggt	taggacgagt	ttaatatagt	gtataaggaa	2160
tttgaaagat	ttaggatattg	tgttttaatt	aacgttaagt	agaatggata	agtttttagt	2220
attttgaaaa	cgttgggtta	gggttttttt	tttattgtgt	gttttttgtt	tggggattaa	2280

taagtattat	agagaacgtg	at ttgagggc	at ttttttatt	tttgtataaa	tttagagtga	2340
attattaaat	agttgttcgt	ttaaagttaa	ggtaattttt	ttttgacggg	tttatttggt	2400
tttcgatttt	taatttatta	gtttgttttt	ttagggtttt	gttttttttg	taattaaagt	2460
tttttttagat	tagcgtagta	tttatttgat	aggttgtttg	gaaaatttaa	gatcggagag	2520
gtgatttggt	gttggttttt	aaatttttta	gttttaagta	acgtgttttt	tttttatatg	2580
gggtggggga	ttggaaatgg	atgtagtgag	atataaagag	tgggtgtttt	gttgattttt	2640
gtattttttt	ttttttgatt	at tttatttt	ttttttttta	gttttcgatt	tttagtttta	2700
tttttttatt	tttggtttcg	tattaaaagt	cggatcgttt	tgggttgggt	aggagttaga	2760
ttttcgggag	tttggtttgtg	tagatttagt	gcgtacggcg	aggtagtagt	tcggtttcgt	2820
attgttgata	ggtgtaggta	ggatagtttt	tttatcgcg	ttcggggcgt	tttgattggg	2880
cgggagttac	gttagtcgta	ttcggagaag	ggtttgggag	gaggcggagg	cggagagggt	2940
tggtaggggt	cgcggcggag	tgacgtttcg	gtattaggaa	gttcgttttt	ggttttaaga	3000
tgttaggtta	atagggaagc	gcggagtcgt	agatttgggt	cgtcgttcgt	ttgggtgttt	3060
ggagttagag	tgcggttaagg	ttcgggtttt	gttcgatcgt	tcgaggggtg	tgctgtgctg	3120
cgttgccggag	ggtgcgttta	gagggtcgcg	tcgtggttgt	agcgttgttt	gtcgtcgtag	3180
gggattttaat	attattttatt	tggttttggt	at ttttgata	tttttttggt	aggggtgtcg	3240
cgtggggggg	gggcgggtag	agcgcggtcg	gcgttagttt	tttttattgg	aggggttttt	3300
gggggaggga	gggagagaag	aagggggttt	ttgtttattt	ttgtttcgtt	ttggagttag	3360
gaagtttggt	ttttaaaagac	gttttgagt	gtgttttttt	gttttatatt	tatgttttcg	3420
tttggttcgt	gatttttcgt	tttcggattt	tttcgtttga	gtttttcggg	ggagacgggg	3480
gtagtttggt	ttgagaattc	ggcgggggtt	gcgttttttg	gtttttttcg	tacgcgggaa	3540
at ttcgcgtt	tagagcgcga	ttcggagcgg	gtagcggcgg	ttacgggggt	tcggcggggg	3600
agtagttaag	gattagtaga	gcgtcgcgtt	ttttcgttta	tgaattgtat	gaaagggtcg	3660
ttttatttg	agtatcgagt	agcggggatt	aagtgttcgg	tcgttttttt	at ttttttgt	3720
tattattttt	agtcgttagt	tatggtttcg	gttttggttt	tcggttagtt	tcggtcgttg	3780
gattttttta	agtataggtt	ggaggtgtat	attattttcg	atatttttag	ttcggagggtc	3840
gtaggtaagg	cgtcgcgtcg	ttttgtagat	at ttttcgtt	agttgttttg	cgttattcgt	3900
tttttttcgt	tttaagggaag	ttagtttttt	cggggggagg	cgtggtggga	gtggtcgttc	3960
gtttggtttt	tcgtagaatt	ttcgggagtc	ggaattttga	ttatttcgta	tttttttagt	4020
tttttttcga	tcggttcggt	ttttggggcg	ttaagggcgc	gagtaatttt	tcgttttttt	4080
ttattcgtat	tttggttttt	tttttggttt	ttgggttata	aaaatttttag	tattttgatt	4140
cgaggatttt	tagagggtcgt	cgatttttgt	ttttgttttt	ttttcgggtt	ttagttttcg	4200
aggagtttta	ttcgttagga	aattgtttga	aattatttag	aatgtttttt	cgcgaagagg	4260
tatttttttt	tttttttttg	gaaagggtcg	gcgaatttcg	gtgtttaatc	gaatttttat	4320
at tttttttt	agttttttta	aatcgtatgg	aaatttgagt	tttttgcgag	ggggaggggg	4380
gtttgtaaat	tacgcgcgtg	tgccgcgttt	aggagatttg	gtgtgtttgc	gtagagggtg	4440
ataaatatat	ttgaaagtat	aggttataaa	agtgaatgtg	tcgtttgtagt	gagataaata	4500
tgtaaaataaa	acgtgcggcg	ttgggggagg	ggaggaaatg	gggcgcggat	at tttatttt	4560
gcgtttgtat	at ttttatagg	cgtagcgttt	ttcgcggttc	ggagtcgtcg	cgcgtatttt	4620
ttttcggcgt	taggtagttt	agttttttta	cgttttttgt	cgtcggttta	gttgccgttc	4680
gcgttgtagg	tgggtatggt	gacgggaaag	tgtgtgtgtt	tcgttttttag	agaaagataa	4740
aagttagtag	gggaagaatg	aggacgtggg	cgtcgaggat	tcgtttaaga	agaagcggta	4800
aaggcggtag	cggattttatt	ttattagtta	gtagtttttag	gagttggagg	ttatttttta	4860
gaggaatcgt	tattcggata	tgttttatac	cgaagaaatc	gttgtgtgga	ttaatatttac	4920
ggaagttcga	gttcgggtag	gagtttagtac	ggagtttggt	agggatgggg	ggaggatggt	4980
gtggaggtag	aggttaagta	gattaggaga	gaatgtggaa	ggtagcgtcg	tttgggaggg	5040
cgtcgggtggg	gcgtagtttt	gtaaaaggtag	aaggttttcg	ggcgggtttg	ttgcgagatt	5100
atagtttttt	tttcgagggtc	gataggattg	tcgttttggt	ttagggtttt	agagcgggat	5160
cggttttattg	tttcgtttatt	tcgcgatttt	acgagttggg	ttgtatgggt	aattttttgt	5220
ataggatatt	gtgttttttg	tttgtagttg	ttagagtaga	gttaataaaa	tttttatttag	5280
gttaagagtc	gcgaatagg	tttaatttgt	gagtttttaa	taaggaaaaat	tcgttagaga	5340
tacggaagag	ttggtttttt	ttgggaaatt	tttggttcgg	ttttggttta	gttttttttt	5400
ttttgggttc	gcgtttttta	tatttttttt	acggttggtt	cggttatttta	ggtttttttt	5460
atataatttta	tttttttagt	ttgtgatttt	cgggagtaaa	gtttttaatat	ataattatta	5520
gttttttttag	aaggagaaag	aaaaaaagaa	gaaagatttt	tttggttggt	ttattttatt	5580
tttttttagga	gttgaatttt	ggaaattgaa	at tttatttt	ttttttttta	attataattt	5640
tagtttttgta	aaaagggttt	at ttttaatt	tgtagtaaat	ttgtatttta	tggattggtta	5700
aaaatgagtt	taaataaata	at ttttaatt	aacgtttttg	tttatgttgg	tcggtggaag	5760
at ttttaaat	tgtaggatt	ttggaagtag	aaaatagaat	taagtaaat	aagcggatt	5820
tagagggttt	gttggttaaaa	aaaaaaaatt	aagtgttttg	ggtagaaaaa	ataaagtttt	5880
cggttagagt	agagtaaaata	aaaagaagaa	aataacgata	aaaagaataa	agattaaaaat	5940
gttttttttaa	at tagaggga	atgaagatat	tttttggttg	gtattttgtg	aaggtagag	6000
gttatgttg	tggataaaaag	gtcgggaaga	agttgaaaat	ggtttttagt	taattgttta	6060

gagttagagt	tgggttttgg	gcggcgtggt	tttgagtaag	gttagttttt	tattagtttt	6120
tttgtatatt	aagggaacgg	gtttttttacg	tatttttttc	gtttgagtaa	agtttagatg	6180
gtttagggta	gaaatggtaa	gtaattaaag	atagagttaa	tgggtttttt	gggatttttc	6240
gaaaacgttt	ttttatttcg	ttcgttattt	cgtagttaa	ttttagtgtt	ttgtagtcgc	6300
ggcgttggtt	tttttttgta	gttgtttttt	tttttagggc	ggttgtttgt	cgagttaagt	6360
gggagtggag	cgtgtttttt	atagtagtcg	ggtgtaaaga	ggaaggggga	taaaaaggaa	6420
attaagaatg	aaaggaaaaa	gagaaaaagc	ggattatacg	gttgggttcg	gcggagatgt	6480
gtaatgtgaa	atattattgg	tgtagtttcg	gatattttag	gttaggtttt	tttttaatat	6540
ataaaagtcg	tcgtttgggg	cgatagggag	gttcgatgtg	gattgggacg	ggggttcg	6600
ttgggttatc	ggatacgggt	ggaagtcggt	cggtttgggt	ggtcgtttgt	aaagttaaac	6660
gattcgtttg	ggtttggcgc	gcggataggt	ttgtggtggg	tttagggtaa	agaagaggta	6720
gagcgaaaga	agggggaatt	tttaaaatta	tttttttcgg	gttttcggag	tttaatatgt	6780
taagtttttg	gagttaacga	gttgacgaag	aggtggtttt	ttgtttttta	tttgggtgtt	6840
ttgttaggcg	agaaagagtg	ttggcgggtt	agtttttggt	aaggaggtac	gtattagggg	6900
gtgggggacg	atagtggagg	ttagggaagg	aaggaggaa	ttgcgtggga	gaaagagcga	6960
tttttttagt	tttttttagt	tttttttttt	tattcgtggg	tttgtggttt	tggaaatggaa	7020
gtaagtttgt	aagggtgttt	gggaagggtt	ggaaaagttt	gttgtttcgc	gtttgtttta	7080
tattaagtgt	ttttggattt	ggagaaacgt	ttggttgagt	gattaaatcg	ttcgtaggtt	7140
tttatgcgtt	cggttgaggt	ttgtggcgta	gtttcgagtt	tttagttcgt	ggttagagta	7200
gattaggttt	tttgcgtttg	gtggagattc	gggttagtaa	ttgaaagttg	gttttggtat	7260
tttggtgtgt	agggcgggtg	agtgaagcga	ggttagggtg	tgtgagtgcg	ttagcgtgtg	7320
tgtcggggga	agggcggggg	tggttttcga	tggaagtttt	agtaatttgt	attgtggtat	7380
ttgtttgttt	ttttgtttta	atcgttttta	ggtttggttt	aagaatcgtc	gggttaaattg	7440
gagaaagagg	gagcgttaatt	agtaggtcga	gttatgtaag	aatggtttcg	ggtcgtagtt	7500
taatgggttt	atgtagtttt	acgacgat	gtatttaggt	tattttttata	ataattgggt	7560
cgttaagggt	tttatattcg	tttttttatt	tattaagagt	tttttttttt	tttaattttat	7620
gaacgttaatt	tttttggtat	tatagagtat	gtttttttta	tttaattttta	tttagttttat	7680
gagtatgtcg	tttagtatgg	tgtttttagt	agtgataggc	gtttcgggtt	ttagtttttaa	7740
tagtttgaat	aatttgaata	atttgagtag	ttcgtcgttg	aatttcgcgg	tgtcgcacgtt	7800
tgtttgtttt	tacgcgtcgt	cgattttttc	gtatgtttat	agggatacgt	gtaattcgag	7860
tttggttagt	ttgagattga	aagtaaagta	gtatttttagt	ttcggttacg	ttagcgtgta	7920
gaattcgggt	tttaatttga	gtgtttgtta	gtatgtagtg	gatcggttcg	tgtgagtcgt	7980
atttatagcg	tcgggatttt	aggattttgt	cggatggggg	aatttcgttt	ttgaaagatt	8040
gggaattatg	ttagaagggtc	gtgggtatta	aagaaaggga	gagaaagaga	agttatatag	8100
agaaaaggaa	attattgaat	taaagagaga	gtttttttga	ttttaaaggg	atgttttttag	8160
tgtttgatatt	tttttattat	aagtattttt	aatagttgta	aggatatata	tataaataaa	8220
tgtttgattg	gatatgatatt	tttaattatta	ttataagttt	gttattttttt	aagttttagta	8280
ttgttaatat	ttaaatgatt	gaaaggatgt	atatatatcg	aaatgttaaa	tttaattttat	8340
aaaagtagtt	gttagtaata	ttataatagt	gttttttaaag	gttaggtttt	aaaataaagt	8400
atgttatata	gaagcgatta	ggattttttc	tttgcgagta	agggagtgt	tatatataat	8460
gttatattgt	atgttttttaa	tatattatta	ttattataaa	aaatgtgtga	atattagttt	8520
tagaatagtt	tttttggtgg	atgtaatgat	gtttttgaaa	ttgttatgta	taattttattt	8580
tgtgtataat	atttcgtata	atattattgt	tttattttttt	agtaaataatg	aaataaatgt	8640
gttttatattt	atgggagtaa	aatatattgt	atataaattg	gttttgattt	ttttttttttt	8700
tttttggtat	taatttggtt	aggatatttt	gtttattgtt	ttttaaataa	attagttttt	8760
tttggttgtt	tagttaaata	tataaggtag	tagttttttat	ttaaatttgg	tagaaataaa	8820
tgatagttat	ttattagaaa	ttaaaaagaa	aaaaaaagggt	attttcgggg	gggaaaaggg	8880
ttataaaatt	taattttgtt	tttttaattt	tttttttggtt	taaatttaga	ggatttttatt	8940
atggttagta	aataatatga	aaaagaaaaa	agaagaaaga	aatttagtaa	gtttatttagt	9000
t						9001

<210> 4

<211> 9001

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 4

agttgatgga	tttggttaaat	ttttttttttt	ttttttttttt	tttatattat	ttgttagtta	60
taatggaatt	tttttaggttt	aagttaaaga	aaaattggag	agataaaaatt	agatttttga	120

gttttttttt	tttttgggaa	tgtttttttt	tttttttttt	gtttttgatg	aatgggttatt	180
atattttttt	attaaattta	aataaggatt	gttgttttgt	atgtttaatt	aggtaggtag	240
aggggaattgg	tttgttttagg	aagtagtgat	tgagatgttt	tggttaagtt	agtगतगग	300
gaggggagaa	agaatttaga	ttaatttgta	tgtagtatat	tttattttta	tgaaataaaa	360
tatattttgtt	ttatatttgt	tgaaaagtaa	aataataata	ttgtatgaaa	tgttatatat	420
agggtagggt	gtatatagta	gttttagaaa	tattattgta	tttatttagag	aaattattttt	480
aaaattgata	tttatatatt	ttttataata	ataataatat	gttagaaata	tatagtgtgg	540
tatttagtat	atataattttt	ttgtttgtaa	gtgaaaaatt	ttaattgttt	ttgtataata	600
tgttttat	ttaaagttaa	tttttaaaaa	tattgttgtg	atattattaa	taattgtttt	660
tataaaatta	atttgatatt	ttgatataata	tatatatttt	tagttattta	aatgttaata	720
atgttaaaatt	taaaaaataa	taagtttata	gtaattgtta	aatgttatat	ttagttaaat	780
atgtgttgt	gtatgtgttt	ttgtaattgt	tagaaatatt	tgtagtgaaa	gatgttagat	840
attgaggata	tttttttgaa	attaaaggag	tttttttttt	gatttagtgg	tttttttttt	900
tttatatagt	tttttttttt	tttttttttt	ttagtgttta	tgatttttta	gtataattttt	960
tagtttttta	agggtggagt	tgttttat	ggtaagggtt	taggattttg	gtgttgtggg	1020
tggtgtttat	atgggttggt	ttattgtata	ttggtaagta	tttaggttgg	agggtgggtt	1080
ttgtatgttg	gtgtagttga	agttggagtg	ttgttttgtt	tttagtttta	ggttgggttag	1140
gtttgagtta	tatgtgtttt	tataaatata	tggaggagt	ggtggtgtgt	aaggataggt	1200
aggtgttgg	attgtggaat	ttagtgtagg	gttatttagg	ttgtttaagt	tatttaggtt	1260
gttgagattg	gagtttggga	tgtttgttat	tggtgagggt	attatgttgg	atgatattgtt	1320
tatggatgag	atagagttgg	gtggggaaaa	tatgttttgt	gatgatagg	ggttgatgtt	1380
tatagagttg	aagaagggga	agtttttgtt	ggataggagg	gtggatgtaa	ggtttttgtt	1440
ggtttagttg	ttgtaggaat	agtttgggta	tatgttgttg	tagggttgta	tgagttttatt	1500
gaattgtggt	ttgaagttat	ttttgtatag	tttggttgtt	tggttgtgtt	tttttttttt	1560
ttatttgggt	tgatgatttt	tgaattaaat	ttgggggtgg	ttggggtaag	ggagtaaata	1620
gatgttatag	tgtagattat	taaaattttt	attggagggt	aatttttgtt	tttttttgat	1680
atatagtta	gtgtatttat	atatttttgt	tttgttttat	tgtattgttt	tgtatatata	1740
gatattaggg	ttagttttta	gttattgggt	tgggttttta	tttaagttag	gagatttggt	1800
ttgttttgg	ttgtgagttg	ggatttggag	ttatgttata	aatttttagt	gaatgtatgg	1860
agatttgtgg	atggtttgat	tatttagtta	ggtgtttttt	taggttttaa	aatatttaatt	1920
gtaaaataaa	tgtggggtag	taggtttttt	taattttttt	tgggggtatt	tgtaaaattt	1980
tttttat	aaagttatag	atattatggat	gaggagaagg	ggttggaagg	gtattagagg	2040
attgtttttt	tttttatgta	atattttttt	tttttttttg	atattttatt	ttgtttttta	2100
ttttttggta	tgtgtttttt	taatagggat	taggttggtta	atattttttt	ttgttttagta	2160
aaataattaa	ataaagagta	aaagattatt	tttttggttag	tttgttaatt	ttaggagt	2220
ggatatattaa	attttgggaa	tttggaagg	gtagttttgg	agattttttt	tttttttgtt	2280
ttgttttttt	tttattttaa	gtttattata	ggtttgtttg	tgtgttaggt	ttagttgggt	2340
tgtttgggtt	tgtaggtgg	tatttaggtt	ggttgggttt	tatttgtgtt	tggtgggtta	2400
gttgtaattt	tgattttta	ttatatggg	tttttttgtt	gttttagatg	gtggtttttg	2460
tgtattggag	agaggtttgg	tttgagatat	ttgagttgat	attagtgatg	tttatatta	2520
tatatattttg	ttgggttttag	ttgtgtaatt	tgtttttttt	tttttttttt	ttatttttga	2580
tttttttttt	atattttttt	tttttttgtat	ttgattgtta	taaaaagtat	gttttat	2640
tatttgggtt	gataagtagt	tgttttggaa	ggagaggtag	ttgtaaggag	agtttagtgt	2700
tggtgtata	aagtattagg	gtggagtgt	ggaatagtgg	gtgggggtgg	agggtgtttt	2760
tgaaggattt	tagaaaattt	atagattttg	tttttaatta	tttgttattt	ttattttagg	2820
ttatttaaat	tttgttttagg	tgagaagagt	atgtgagagg	tttgtttttt	tgatgtgtaa	2880
gagagtta	gaaagattga	ttttgtttta	aattatgttg	tttaggattt	agttttgggt	2940
ttggatagtt	aaattaaaat	tatttttta	ttttttttgg	ttttttattt	attagtata	3000
ttttatgttt	tgtataaatg	ttatttagag	agtgttttta	ttttttttga	tttgggagag	3060
tattttgggt	tttatttttt	ttattgttgt	tttttttttt	ttgtttgttt	tgtttttaatt	3120
gggggtttta	ttttttttat	tttagagtatt	taattttttt	tttttaatat	taaagttttt	3180
ggatgttgtt	tgatttgttt	gattttgttt	tttgttttta	gaattttta	aaatttggaa	3240
ttttttattg	attagtataa	attaggatgt	tgttattggg	ttattttatt	gagtttat	3300
ttgttaattt	ataaagtata	gatttgttat	aaagttaagg	taagtttttt	ttataaaatt	3360
atgattataa	tttagaagag	gggtgtgtag	ttttaatttt	tagagtttaa	tttttgagag	3420
aagataaata	aattaaagtag	aaaagttttt	tttttttttt	tttttttttt	tttaagagga	3480
ttagtagttg	tgtattaaaa	ttttgttttt	ggagattata	aaattaggaa	atagggtgtg	3540
tgggagagat	ttgaatgggt	gaaataattg	taaagaaggt	gtaagaagtg	tgagtttagg	3600
agggaaaaag	ttgggttagg	gttgggataa	agggttttta	gggagggtta	atttttttgt	3660
gtttttgggt	ggtttttttt	gttaaagggt	tatagggttg	agtttgtttg	tggttttttg	3720
tttggtagg	attttatttag	ttttgttttg	gtaattgtta	gttaggaata	taattgtttg	3780
tgtaggggat	tgtttatgta	gttttagttg	tgagattgtg	ggatgggtgg	gtagttaggt	3840
ggtgttgttt	tgggagtttg	agtttaggtg	gtagtttgtt	tggttttgg	gagggaaattg	3900

taatstttgtat	attaggtttgt	tgtgaggtttt	tttgttttttg	taaagtttgtg	ttttatttggt	3960
gttttttttag	gtggtgtttgt	tttttatattt	ttttttttggt	ttattttgggt	tgtatttttta	4020
taatatstttt	tttttatsttt	tttttagattt	tgtgtttgggt	tttattttgga	tttgggttttt	4080
tgtataggttg	gttttatatag	tgatttttttt	gtgtgttggt	atgttttgggt	agtgggttttt	4140
ttggaaagtgt	gttttttagtt	tttggagtttg	ttgggttggtat	aagtgtagttt	gttgtttgttt	4200
ttgtttgtttt	tttttagatgt	ggttttttgggt	gttttatgttt	ttatttttttt	tttgtttgggt	4260
ttttatttttt	tttgaaaatgt	aaatatatat	attttttttgt	tagtatgtttt	atttgtaatgt	4320
tggatgttaa	ttggatttgggt	ggtagaagtt	gtggaagagt	tgggtttgttt	ggtgttggtg	4380
gaggtgtgtgt	gtggttgggtt	tgggtttgtga	ggagtgtttgt	gtttgtgggtt	tgtgttaggtg	4440
taagtgtgggt	tgtttgtgttt	ttatttttttt	tttttttttta	gtgtttgtatg	ttttattttat	4500
atgtttatttt	tattgtagtgt	gtatatstttat	tttttatagtt	tgtgtttttta	agtatatsttta	4560
tatatsttttt	tgtatagatata	ttaaatttttt	tgggtatgtgt	atatgtgtgtgt	ggtttataga	4620
ttttttttttt	ttttgtagaa	agttttagatt	tttatgtgtgt	ttgggaaggt	taggaaaaga	4680
tgtgggtgatt	tgggttgggtat	ttgaagttttg	ttgggtttttt	tttaaaaaaa	aaaaaaaat	4740
gttttttttgt	gaaggttatt	tttgagtgtgt	tttaggtaat	tttttaataga	gtggagttttt	4800
ttgggtgattg	aaagtgtgaga	ggaaaatagg	gatagaggtt	ggtgtgttttt	gaaggtttttt	4860
gaattaagat	gttgggtattt	ttgtgatttta	ggaaatagaa	gggaggttag	ggtatgaata	4920
gagaggtgtg	tagaattgttt	tgtgtttttta	gtgtttttagg	agtgtgggtt	gttgaggtgag	4980
aattaaggtg	atgttgggtat	gttaaaaattt	tgggttttttg	aagtttttgtg	gggaggttagg	5040
tgaatgattat	ttttttattat	gtttttttttt	ggaggtgtgtg	atttttttttg	ggtgagaggtg	5100
agtgggtgtgt	gtatagtagtt	tgtgtgggtat	tgtttgttagg	gtgtgtgtgtgt	gtttttatttt	5160
tgggttttttg	gttggaggtgt	ttggagatgtg	tgtgtattttt	tagttttgtgt	ttggaggtggt	5220
ttagtgtattg	gggtttgattg	ggagttagaa	ttgaagttat	ggttaattgtt	tgggtgattgt	5280
gataggaaga	ttaggtgattg	gttgatagtt	tgggtttttgt	tgtttgtgtgt	tttaagttaa	5340
gtgggtttttt	tattgtagttt	atggatgagg	gagtgtgatg	ttttattagt	ttttgtttat	5400
tgtttttgttg	agtttttgtat	gttgtttgttg	tttgttttttg	gttgtgttttt	aggtgtgggtg	5460
ttttttttgtt	gtgggtgagag	ttaggtgtgt	taattttttgt	ttagtttttta	agttaagtgtg	5520
ttttttgtttt	ttttggaaag	tttaagttaa	aaagttttga	gatggaaagt	tagtgggttaa	5580
atgaagatat	gggtatgtgtg	tagaaggtgt	ttatttttag	tgttttttag	gagttaggttt	5640
ttatgttttta	aagtgaataa	agagtgggtat	aagattttttt	ttttttttttt	ttttttttttt	5700
taagaattttt	tttaataagg	aaagttaattg	ttgattgtgt	tttgtttgttt	tttttttttat	5760
gtggttagttt	ttagatagaa	gtgttaagag	ttagatgggt	aggttaggtga	tatttagattt	5820
tttgtgtgtg	tagtagattgt	tgttagttatg	atgtgtgtttt	ttgagtgtat	ttttttgtaat	5880
gtgtatatgt	atatttttttg	ggtgtttttaa	taggtagtttg	gtttttgtttg	agtttagtttt	5940
taggtatttta	ggtgagtgtat	ggatttagatt	tgtgttttttg	tgttttttttg	tttgttttaatt	6000
atttttaaaat	tagaggtgtg	ttttttgtgt	ttgagattgtt	attttgtttgt	ggttttttttt	6060
agttttttttt	gtttttgtttt	tttttttagat	tttttttttg	gtgtgattga	tgtgtgttttg	6120
tattatattag	gatgtttttga	gttgtgtgtg	aggtattgttt	ttgtttgtat	ttatttagtag	6180
tgtgtgtgttg	ggttattgttt	ttgtttgtgtg	tattgtgtttt	atataggttaa	gttttttggga	6240
attttagtttt	tgttttagttt	aaggtgatttt	ggttttttagt	atgaatttaa	aggtgaagag	6300
atgaggttag	gagtgtgaag	tttgggtgaa	gagagtggaa	tgggttaagaa	gagaaaggtat	6360
taaggatttaa	taagatatatt	atttttttgtg	ttttattata	tttatttttta	attttttatt	6420
ttatatataaa	aggagatatg	ttattttaaa	ttgaaaaatt	tgaaaaaatag	taataaaatta	6480
ttttttttgat	tttaaatsttt	ttaaatagtt	tgttaagtga	atgtttgtgtt	aatttgaaga	6540
agtttttaatt	gtaaaagaaga	tagagtttttg	aaaaggttag	ttatataaatt	agaaatttag	6600
aagtaaatgt	atttgtttaa	agaaaattat	tttgattttta	aatgaataat	tgtttgtgtg	6660
tttatsttttg	atttatataa	gaataaaaaag	ttgtttttaga	ttatgtttttt	tgtgatgtttt	6720
attagtttttt	agatagaaaa	tatatataatg	aagagaaatt	ttatatttagt	gttttttaaaa	6780
tgttgaaagt	ttattttattt	tatttaattgt	tgtttaagat	atatattttta	gatttttttaa	6840
attttttgtat	tattgtattat	agtttgttttt	aatttgagag	agttatgtttt	taaattttgat	6900
ttttttgtttt	atttttattat	taatttagatt	taaattttata	aagtttttag	aattaataat	6960
tttgagtttaa	ttatatatga	aatatgttttt	aatgaattttt	tatatataatg	agaattgtgt	7020
taaataatta	atttttaagga	taattttttta	tagttattttt	tttttttttag	ttagttttaag	7080
ggtgtttttga	gtttattaaag	tttaagttag	tagaaggtgt	gtgtgtgtagt	taaggtgtgaa	7140
aagtttttagaa	ttgtgttttaa	tttagtaaaag	taaaattttta	tttatataaaa	ataaaaaaaa	7200
ttatttttttg	agatatataat	tttttatagtt	attgtttttta	agtaaaattta	attttttaaag	7260
aaattaaaga	aagaaatttta	aatatatttta	aaataattttt	tgaaggttttt	tttgttttttt	7320
agtataggttt	agttggagag	gataaaattaa	tttttttttgg	gttttttgtat	gggtgatttgt	7380
tttattatgtg	agtttagtgtt	attattttttg	aatgtgtattt	tgtttgatatt	tatagtttaatt	7440
gattttgtaatt	gttagtatga	agtattttttta	aaatatstttt	ttttttgtttt	tgttttataag	7500
atttggaaatt	ttattttgatg	tggaaataaag	tggatgaagt	agattataaaa	tatatatttga	7560
atttatgtgtt	ttttttttttg	ttttgtattat	tttttaaatstt	tattttgtaat	tttttttttat	7620
tttaaatsttg	tagttttaaag	atgtatatga	gaattgttttt	ttagttttttt	tttatttagta	7680

ttatTTTTatt	ttaagaataa	tttagttgta	agggaggaat	ttttttatag	taagtttttaa	7740
attagTatttt	ttgttttttaa	ttttttatttt	tatttttattt	tatttttatat	atatagatat	7800
ttgttttagag	taaaatatat	ttttatgtga	taggtttgta	ttagttgagg	tttatatat	7860
tagttatatt	aggttttgta	attttattat	taaattatat	atattatatt	agtagtttgt	7920
tggtaaagaa	ggttaaatta	atttatatttt	tgtttattat	ttgggtgttta	aatgatgtat	7980
tttatttttg	agattttggtg	gagaatttttt	tttttagatt	ttatagtgtt	ttattgaaga	8040
taatgttttt	atattttgtag	tggttttttaa	tttgataaga	ttttaatttg	tttaagtttt	8100
ttaaataagg	gtttttaaatg	tttttagttg	tttttttatt	gaattttttt	taattttttt	8160
aagattataa	agtatatgtg	taaagtaaatt	attttttttt	attgtattgt	tagttgatga	8220
tttataatta	agtttaataag	aatttagttt	ttttttgttg	aatgtgttta	tttaattatat	8280
tttagttttt	ttttttaaat	tttgaataag	ttgtggtttt	tataatatta	tgtttttttaa	8340
agttttattt	tatgaaggga	ttttattata	ttaaagaattg	aaaaaaattt	ttattgtagt	8400
tagtatatat	agttttttat	tttttgtttt	tttaagattta	aatttttagag	ttgtaaatat	8460
ttttggaagt	ttgggtgtta	atgtttttat	ttagaaagtt	gagaagtttt	atagagttat	8520
atagattttt	aaattttatt	tttataaatt	tatagaattt	tgataaaaagt	tttgggtggt	8580
ttattttatt	gatggaattt	ttattatgat	aaatatatat	gtatgaaggga	ttttaatttag	8640
tttttaaagt	ggttgaaaaa	tttaagggtta	tgtgattgtt	ttttatagt	ttaatgtgtg	8700
tgagatgttg	gaagtattgg	ggattagtag	tagtttagat	gtttaaaaag	ataagggtgt	8760
ttaatttggtg	tggattttat	gaagttaagt	ggtgaataaa	gataattatt	tagataattt	8820
agattaaagt	aaaagtaaaa	ttatattttat	ttgtatatat	atattttatat	ttatttttata	8880
ttatagatat	atatatgtat	atatatattg	gtttttgtaaa	taattgattt	aaagtgagga	8940
ttttttttgt	attttttttag	taggagtttt	aatattttttt	taattttttta	attatttttat	9000
a						9001

<210> 5

<211> 9001

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 5

tgtaaagtga	ttaggagatt	aggagaatgt	tgaaatTTTT	gttggaaaaa	tgtaaagaaa	60
atTTTTattt	tgagttagtt	gtttatagag	ttagtgtgtg	tgtgtgtgtg	tgtgtttgta	120
atataaaatg	gatgtgaata	tatatatata	aatagatatg	gttttgTTTT	tatttttaatt	180
tgaattattt	agataattgt	ttttattttat	tatttgattt	taatgggttt	atataaatta	240
ggataatttta	tttttttagg	tatttaggtt	gttggtgatt	tttagtgttt	ttaatatttt	300
gtatatgttg	gtattatgag	gagtagttat	gtgttttttg	gttttttaatt	tattttggag	360
gttgattgag	gttttttata	tatgtatatt	tgttgtgatg	aaagttttat	tggtagagtg	420
gagttatttag	agttttttatt	aaaatttttgt	gggttttatga	gagatgggtt	tagaaattta	480
tatggtttttg	tggggttttt	tgggtttttta	aaataagggt	ttaatattta	agttttttaa	540
aatattttgta	gttttggggt	ttgaattttg	aaaaataagg	agtgaagggt	tgtgtatatt	600
aatattagtg	gagatttttt	ttattttttta	atgtgatgga	gtttttttat	gaaatgaagt	660
tttaaggggt	atggtattgt	ggggattata	gttattttga	ggtttaaaag	aagaaattgg	720
aatatgatta	gtaaatatat	ttagttagaaa	agagttggat	ttttattgat	ttagttagat	780
gttattgggt	ggtagtgtaa	tgggaggaaa	tattttattt	atatatatat	tttatgattt	840
tgggggaatt	agaggaaatt	taataagaaa	atggttagaa	atattttaaaa	tttttattta	900
aaagatttaa	gtaaattaga	gtttttattag	attaaaaatt	attataaatg	taagagtatt	960
gttttttagtg	aaatgtttgt	gggtttgaga	aggagatttt	ttgttaaatt	tttgggataa	1020
aatgtgttat	tttaagtatta	gataatgagt	agaatgtaaa	ttaatttaat	tttttttatt	1080
aataggttgt	tagtgtaatt	tgtataaatt	agtataaga	ttgtaggatt	taatatagtt	1140
ggatgtatga	gttttagtta	atgtagattt	gttatatgag	gatgtgtttt	attttgagta	1200
gggtgtttgta	tgtgtggaat	ggggtaaagt	ggaataaaaag	gttaaaaagta	gaaatgttga	1260
tttaaggttt	attatgaaga	aatttttttt	ttgtagttaa	attattttta	aagtgggatg	1320
atattgggtga	agaaagattg	aaaaataatt	tttatgtgtg	tttttggttt	gtaagtttaa	1380
aatggggagg	agttgttagat	agggttttggg	ggtgggttagg	gtaaaggaga	gatataaag	1440
ttgtaaatat	atttgtagtt	tgtttttattt	attttgTTTT	atattgaata	agttttttta	1500
ttttgtgaat	aaggataagg	agggagtgtt	ttaaagatat	tttatgtttg	tattgtaaat	1560
tattgattgt	aatgttaaat	aaatatatat	ttagagatga	taatatataat	tttatagtaa	1620
aataattgtt	tatgtagaaa	tttagaggag	attagtttgt	tttttttagt	tgattttatgt	1680
tgggggataa	aaggattttt	aaaaattatt	ttgaatatgt	ttggattttt	tttttttaatt	1740

tttttggaaa	ttaaatttgt	ttggaaatag	tgttataaag	agttgatggt	tttaaagggtg	1800
atTTTTTTTT	ttttatataa	ataagggtttt	gtttttgtta	gttgagtgtg	gttttaggtt	1860
ttttgttttt	agtttatata	tatttttttt	gtttgtttgg	atTTtaattgg	tttaagatag	1920
ttttgagttt	attgggaaaa	gaaaatgatt	gttaaaaaatt	atTTTTgaaa	ttggttattt	1980
ggtaatatatt	ttaatgtgat	ggaaatttat	taaggatatat	tttatatata	attagtttaa	2040
ggttgttgat	tttataggtt	ttatggattt	aaatttgatt	gataataaag	taaataagag	2100
agttgaattt	aaagtgtggt	ttttttgggt	taggatgagt	ttaatatagt	gtataaggaa	2160
tttgaaagat	ttaggatatg	tgttttaatt	aatgttaaagt	agaatggata	agtttttagt	2220
atTTTTgaaa	tgttgggtta	gggttttttt	tttattgtgt	gttttttgtt	tggggattaa	2280
taagtattat	agagaatgtg	atTTtgagggtg	atTTTTtatt	tttgtataaa	tttagagtga	2340
attattaaat	agttgtttgt	ttaaagttaa	ggtaattttt	ttttgatggg	tttatttgtt	2400
ttttgatttt	taatttatta	gtttgttttt	ttagggtttt	gttttttttg	taattaaagt	2460
tttttttagat	tagtgtagta	tttatttgat	aggttgtttg	gaaaatttaa	gattggagag	2520
gtgatttgtt	gttgtttttt	aaatttttta	gttttaagta	atgtgttttt	tttttatatg	2580
gggtggggga	ttggaaatgg	atgtagttag	atataaagag	tgggtgtttt	gttgattttt	2640
gtattttttt	ttttttgatt	atTTtatttt	ttttttttaa	gtttttgatt	tttagtttta	2700
tttttttatt	tttgggtttg	tattaaaagt	tggattgttt	tgggttgggt	aggagttgaa	2760
tttttgggag	tttgtttgtg	tagatttagt	gtgtatgggtg	aggtagtagt	ttggttttgt	2820
attgttgata	ggtgtaggta	ggatagtttt	tttattgtgg	tttgggtgtg	tttgattggt	2880
gtggagttat	gttagttgta	tttggagaag	ggtttgggag	gaggtggagg	tggagagggg	2940
tggggagggt	tgtggtggag	tgatgttttg	gtattaggaa	gtttgttttt	ggttttaaga	3000
tgttaggtta	atagggaagt	gtggagtgtg	agatttgggt	tgttgtttgt	ttgggtgttt	3060
ggagttgagt	tgtggttaagg	tttgggtttt	gtttgattgt	ttgaggggtg	tgtgtgtgtg	3120
tgttgtggag	ggtgtgttta	gagggttgtg	ttgtggttgt	agtgggttgt	gttgttgtag	3180
gggatttaat	attatttatt	tgttttttgt	atTTtttgata	tttttttgtt	aggggttgtt	3240
tgtggggggg	gggtgggttag	agtgtggttg	gtgttagttt	tttttatttg	aggggttttt	3300
gggggaggga	gggagagaag	aagggggttt	ttgtttattt	ttgttttgtt	ttggagtgtg	3360
gaagttttgt	ttttaaagat	gttttgagtg	gtgttttttt	gtttatatatt	tatgtttttg	3420
tttgttttgt	gattttttgt	ttttggattt	ttttgtttga	gttttttggg	ggagatgggg	3480
gtagttttgt	ttgagaattt	ggtgggggtt	gtgttttttg	gttttttttg	tagtggggaa	3540
atTTttgtgt	tagagtgtga	tttggagtgg	gtagtgggtg	ttatgggggt	ttgggtgggt	3600
agtagttaag	gattagtaga	gtgttgtgtt	tttttgttta	tgaattgtat	gaaagggttg	3660
ttttatttgg	agtattgagt	agtggggatt	aagttgttgg	ttgttttttt	atTTtttttgt	3720
tattattttt	agttgttagt	tatggttttg	gttttgggtt	ttgggttagt	ttgggttgtt	3780
gattttttta	agtatagggt	ggaggtgtat	attatttttg	atattttttag	tttgagggtt	3840
gtaggtaagg	tgttgtgttg	ttttgttagat	atTTtttgtt	agttgttttg	tgttatttgt	3900
ttttttttgt	tttaagggaag	ttagtttttt	tggggggagg	tgtggtggga	gtggttgttt	3960
gtttggtttt	ttgtagaatt	tttgggagtt	ggaattttga	ttatttttga	tttttttagt	4020
ttttttttga	ttgggtttgg	ttttgggggtg	ttaaagggtg	gagtaatttt	gttgtttttt	4080
ttatttgtat	tttgggtttt	tttttgtttt	ttgggttata	aaaatttttag	tatttttgatt	4140
tgaggatttt	tagaggttgt	tgatttttgt	tttttgtttt	tttttgggtt	ttagtttttg	4200
aggagtttta	tttgttagga	aattgtttga	aattattttag	aatgtttttt	tgtgaagagg	4260
tatttttttt	tttttttttg	gaaagggttg	gtgaattttg	gtgttttaatt	gaattttttat	4320
atTTtttttt	agttttttta	aattgtatgg	aaatttgagt	tttttgtgag	ggggaggggg	4380
gtttgtaaat	tatgtgtgtg	tgtgtgtttt	aggagatttg	gtgtgtttgt	gtagagggtg	4440
ataaatatat	ttgaaagtat	aggttataaa	agtgaatgtg	ttgttgtagt	gagataaata	4500
tgtaaataaa	atgtgtgggtg	ttggggggagg	ggaggaaatg	gggtgtggat	atTTtatatt	4560
gtgtttgtat	atTTtatagg	tgtagtgttt	tttgtgggtt	ggagttgttg	tgtgtatttt	4620
tttttgggtg	taggtagttt	agttttttta	tgggtttttgt	tgttgggtta	gttgggtgtt	4680
gtgtttagag	tgggtatgtt	gatgggaaag	tgtgtgtgtt	ttgttttttag	agaaagataa	4740
aagtttagtag	gggaagaatg	aggatgtggg	tgttgaggat	ttgtttaaga	agaagtggta	4800
aagggtggtag	tggatttatt	ttattagtta	gtagttttag	gagttggagg	ttatttttta	4860
gaggaattgt	tatttggata	tgtttatagt	tgaagaaatt	gttgtgtgga	ttaatttttat	4920
ggaagtttga	gtttgggttag	gagtttagtat	ggagttttggg	agggatgggg	ggaggatgtt	4980
gtggaggtat	aggttaaagta	gattaggaga	gaatgtggaa	ggtagtgttg	tttgggaggg	5040
tgttgggtgg	gtgtagtttt	gtaaaggtag	aagggtttgt	gggtggtttg	ttgtgagatt	5100
atagtttttt	ttttgagggt	gataggattg	ttgttttggg	ttagggtttt	agagtgggat	5160
tggtttattg	ttttgttatt	ttgtgatttt	atgagttggg	ttgtatgggt	aattttttgt	5220
ataggatatt	gtgttttttg	tttgtagtgt	ttagagtaga	gttaataaaa	tttttattag	5280
gttaagagtt	gtgaatagggt	tttaattttgt	gagtttttta	taaggaaaaat	ttgttagaga	5340
tatggaagag	ttggtttttt	ttgggaaatt	tttgttttgg	ttttgggtta	gttttttttt	5400
ttttgggttt	gtgtttttta	tatttttttt	atgggtgttt	tgggtattta	ggtttttttt	5460
atataatttta	tttttttagtt	ttgtgatttt	tgggagtaaa	gttttaatat	ataattatta	5520

gttttttttag	aaggagaaag	aaaaaaagaa	gaaagatttt	tttgtttggt	ttattttattt	5580
tttttttagga	gttgaatttt	ggaaattgaa	atttatatatt	tttttttttaa	attataatta	5640
tagtttttgta	aaaagggttt	attttaattt	tgtagtaaat	ttgtatttta	tggattggta	5700
aaaatgagtt	taaataaata	atttaatagt	aatgttttg	tttatgttgg	ttggtggaag	5760
attttaaatt	tgttaggatt	ttggaagtag	aaaatagaat	taagtaaaatt	aagtggatt	5820
tagaggtttt	gttggttaaaa	aaaaaaaatt	aagtgttttg	ggtagaaaaa	ataaagtttt	5880
tgggttagagt	agagtaaata	aaaagaagaa	aataatgata	aaaagaataa	agattaaaat	5940
gttttttttaa	attagagggg	atgaagatat	tttttggttg	gtatttgtgt	aagggtatgag	6000
gttatgttgg	tggataaaaag	gttggaagaa	agttgaaaat	ggtttttagtt	taattgttta	6060
gagtttagagt	tgggttttgg	gtggtgtggt	tttgagtaag	gttagttttt	tatttagtttt	6120
tttgtatatt	aagggaatgg	gttttttatg	tatttttttt	gtttgagtaa	agtttagatg	6180
gttttagggta	gaaatggtaa	gtaattaaag	atagagttaa	tgggtttttt	gggatttttt	6240
gaaaatgttt	ttttattttg	tttggtattt	tgtagtttta	ttttagtgtt	ttgtagtgtt	6300
ggtgttgggt	tttttttgta	gttggttttt	tttttagggt	ggtgtttgt	tgagtttaagt	6360
gggagttagg	tgtgtttttt	atagtagttg	ggtgtaagaa	ggaaggggga	taaaaaggaa	6420
attaagaatg	aaaggaaaaa	gagaaaaagt	ggattatatg	gttgggtttg	gtggagatgt	6480
gtaatgtgaa	atattattgg	tgttagtttg	gatattttag	gttaggtttt	tttttaatat	6540
ataaaagtgt	ttgtttgggg	tgatagggag	gtttgatgtg	gattgggatt	ggggttgttg	6600
ttgggttatt	ggatatgggt	ggaagttggt	tggtttgggt	ggttgtttgt	aaagttaaatt	6660
gatttgggtg	ggtttgggtg	gtggataggt	ttgtggtggg	tttagggtaa	agaagaggta	6720
gagtgaagaa	agggggaatt	tttaaaatta	ttttttttgg	gtttttggag	tttaatatgt	6780
taagtttttg	gagttaatga	gttgatgaag	aggtggtttt	ttgtttttta	tttgggtgtt	6840
ttgttaggtg	agaaagagtg	ttggtgggtt	agttttttgt	aagggaagtat	gtattagggg	6900
gtgggggatg	atagtggagg	ttagggaagg	aagggaaggaa	ttgtgtggga	gaaagagtg	6960
tttttttagtg	tttttttagt	tttttttttt	tatttgtggg	tttgtgggtt	tggaaatggaa	7020
gtaagtttgt	aaggtgtttt	gggaagggtt	ggaaaagttt	gttgttttgt	gtttgtttta	7080
tattaagtgt	ttttggattt	ggagaaatgt	ttgggttagt	gattaaattg	tttgtaggtt	7140
tttatgtgtt	tgggttaggt	ttgtggtgta	gttttgagtt	ttagtttgta	ggtagagta	7200
gatttagttt	tttgtgtttg	gtggagattt	gggttagtaa	ttgaaagttg	gttttggtat	7260
tttgggtgtg	aggtggtgtg	agtgaagtga	gggttaggtg	tgtgagtgtg	ttagtgtgtg	7320
tgttggggga	aggtgggggt	tgttttttga	tggaaagttt	agtaatttgt	attgtggtat	7380
ttgtttgttt	ttttgtttta	attgttttta	ggtttgggtt	aagaattggt	gggttaaagt	7440
gagaaagagg	gagtgtaat	agtaggttga	gttatgtaag	aatggttttg	gggttaggtt	7500
taatgggttt	atgtagtttt	atgatgatat	gtatttaggt	tattttttata	ataattgggt	7560
tgttaagggt	tttatatttt	tttttttatt	tattaagagt	tttttttttt	tttaattttat	7620
gaatgttaat	tttttgttat	tatagagtat	gtttttttta	tttaattttta	ttttgtttat	7680
gagtatgttg	tttagtatgg	tgttttttagt	agtgataggt	gttttgggtt	ttagttttaa	7740
tagtttgaat	aatttgaata	atttgagtag	tttgttgttg	aattttgtgg	tgttgatgtt	7800
tgtttgtttt	tatgtgttgt	tgattttttt	gtatgtttat	agggatatgt	gtaatttgag	7860
tttgggttagt	ttgagattga	aagtaaagta	gtatttttagt	tttgggttatg	ttagtgtgta	7920
gaatttgggt	tttaatttga	gtgtttgtta	gtatgtagt	gattgggttg	tgtgagttgt	7980
atttatagtg	ttgggatttt	aggattttgt	tggatggggg	aattttgttt	ttgaaagatt	8040
gggaattatg	ttagaagggt	gtgggtatta	aagaaaggga	gagaaagaga	agttatatag	8100
agaaaaggaa	attattgaat	taaagagaga	gtttttttga	tttttaaagg	atgttttttag	8160
tgtttgatat	tttttattat	aagtattttt	aatagttgta	aggatatata	tataaataaaa	8220
tgttttagatt	gatagatat	tttaatatata	ttataagttt	gttattttttt	aagtttagta	8280
ttgttaaat	ttaaatgatt	gaaaggatgt	atatatatgt	aaatgttaaa	tttaattttat	8340
aaaagtagtt	gttagtaata	ttataatagt	gtttttaaag	gttaggtttt	aaaataaagt	8400
atgttatata	gaagtgatta	ggattttttg	tttgtgagta	agggagtgt	tataatgaat	8460
gttatattgt	atgtttttta	tatatattata	ttattataaa	aaatgtgtga	atattagttt	8520
tagaatagtt	tttttgggtg	atgtaatgat	gtttttgaaa	ttgttatgta	taattttattt	8580
tgtgtataat	attttgtata	atattattgt	tttatttttt	agtaaataatg	aaataaatgt	8640
gttttatttt	atgggagtaa	aatatatatt	atataaattg	gtttggattt	tttttttttt	8700
tttttgggtt	taatttgggt	aggatatatt	agttattgtt	ttttaaataa	attagttttt	8760
tttgggtgtt	tagttaaaaa	tataaggtag	tagttttttat	ttaaatttgg	tagaaataaaa	8820
tgatagttat	ttattagaaa	ttaaaaagaa	aaaaaaaggt	atttttgggg	gggaaaagg	8880
ttataaaatt	taattttgtt	tttttaattt	tttttgggtt	ttaaatttaga	ggattttatt	8940
atgggttagta	aataatatga	aaaagaaaaa	agaagaaaga	aatttagtaa	gtttattagt	9000
t						9001

<210> 6
<211> 22
<212> DNA

<213> Artificial Sequence

<220>

<223> PRIMER

<400> 6

gtaggggagg gaagtagatg tt

22

<210> 7

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> PRIMER

<400> 7

ttctaatacct cctttccaca ataa

24

<210> 8

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PROBE

<400> 8

agtcggagtc gggagagcga

20

<210> 9

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> PROBE

<400> 9

agttggagtt gggagagtga aaggaga

27

<210> 10

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> PRIMER CONTROL

<400> 10

tggtgatgga ggaggttttag taagt

25

<210> 11

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> PRIMER CONTROL

<400> 11

aaccaataaaa acctactcct cccttaa

27

<210> 12

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> PROBE CONTROL

<400> 12

accaccacccc aacacacaat aacaaacaca

30

<210> 13

<211> 408

<212> DNA

<213> Homo Sapiens

<400> 13

tcctcaactc	tgcaaggcctg	aaagaaggtc	acacacgcac	gctcacaccc	acactccaca	60
cgctcgtcc	caaacaaccc	catgaacatt	gtcctttgtt	ccgtctcttg	ggccactttc	120
cctgtcgctt	cctcccagcc	cgctctgatt	tgctcccca	aagtacgttt	ctgtctcccc	180
gctgcccctg	cgctcccct	ttgatttatt	agggtgccc	ggttggcgca	gattgctttt	240
tcttctcttc	catcccatcc	tcccttcttg	tctcctttc	cacagtggga	gtccgtgctc	300
ctgctcctcg	gttggtcct	aagtgcccc	ccaggtcccc	tctcctttcg	ctctcccggc	360
tccggctccc	gactcttcgg	cccgtggca	tctgcttccc	tcccctgc		408

<210> 14

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 14

aacatctact tccctcccct ac

22

<210> 15

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 15

gttagtagag attttattaa attttattgt at

32

<210> 16

<211> 14

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 16

ttcgggttgcg cggt

14

<210> 17

<211> 16

<212> DNA

<213> Artificial Sequence

<220>

<223> chemically treated genomic DNA (Homo sapiens)

<400> 17

tttggttggtg tgggtg

16

<210> 18

<211> 144

<212> DNA

<213> Homo Sapiens

<400> 18

ttctggtcct cctttccaca gtgggagtcg gtgctcctgc tctcgggttg gctcctaagt
gccccgccag gtccctctct ctttcgctct cccggctccg gctcccgact cttcggcccc
ctggcatctg cttccctccc ctgc

60

120

144

<210> 19

<211> 162

<212> DNA

<213> Homo Sapiens

<400> 19

tggcatctgc ttccctcccc tgctcgttt ctcgtcgccc ctgctcgctc cccccggcgc
tcgcccgggc gctgtgctcg ctctggatc gccagccgcg cagcgggctc gccggcgccc
gcgcgccact gtgcagtga gtttggtgga atctctgctg ac

60

120

162

<210> 20

<211> 2235

<212> DNA

<213> Homo Sapiens

<400> 20

tgggagtcgc	tgctcctgct	cctcgggttg	ctcctaagtg	ccccgccagg	tccctctctc	60
tttcgctctc	ccggctccgg	ctcccgactc	ttcggcccgc	tggcatctgc	ttccctcccc	120
tgctcgttt	ctcgtcgccc	ctgctcgctc	ccccggcgcc	tcgcccgggc	gctgtgctcg	180
ctcctggatc	gccagccgcg	cagccgggct	cggccggccc	cccgcgcgcc	actgtgcagt	240
ggagtttgg	ggaatctctg	ctgacgtcac	gtcaactccc	acacggagta	ggagcagagg	300
gaagagagag	ggatgagagg	gagggagagg	agagagagtg	cgagaccgag	cgagaaaagt	360
ggagaggagc	agaaagaaac	tgccagtggc	ggctagattt	cggaggcccc	agtgcacccg	420
tggactcctt	cggaaacttg	cacctcagg	agccctgcag	tcctctcagg	cccggctttc	480
gggcgcttgc	cgtgcagccg	gaggctcgcc	tcgctggaaa	tcgcccgggg	aagcagtggg	540
acgcggagac	agcagctctc	tcccggtagc	cgataacggg	gaatggagac	caactgccgc	600
aaactggtgt	cggcgtgtgt	gcaattaggc	gtgcagccgg	cggccgttga	atgtctcttc	660
tccaaagact	ccgaaatcaa	aaaggctcag	ttcacggact	ctcctgagag	ccgaaaagag	720
gcagccagca	gcaagttctt	cccgcggcag	catcctggcg	ccaatgagaa	agataaaaagc	780
cagcagggga	agaatgagga	cgtgggcgce	gaggaccegt	ctaagaagaa	gcccgaagg	840
cggcagcgga	ctcactttac	cagccagcag	ctccaggagc	tggaggccac	tttccagagg	900
aaccgctacc	cggacatgtc	cacacgcgaa	gaaatcgctg	tgtggacca	ccttacggaa	960

gcccagagtc	gggttttggt	caagaatcgt	cgggccaaat	ggagaaagag	ggagcgcaac	1020
cagcaggccg	agctatgcaa	gaatggcttc	gggccgcagt	tcaatgggct	catgcagccc	1080
tacgacgaca	tgtacccagg	ctattcctac	aacaactggg	cgcceaaggg	ccttacatcc	1140
gcctccctat	ccaccaagag	cttccccttc	ttcaactcta	tgaacgtcaa	ccccctgtca	1200
tcacagagca	tgttttcccc	acccaactct	atctcgtcca	tgagcatgtc	gtccagcatg	1260
gtgccctcag	cagtgcacag	cgtcccgggc	tccagtctca	acagcctgaa	taacttgaac	1320
aacctgagta	gcccgtcgct	gaattccgcg	gtgccgacgc	ctgcctgtcc	ttacgcgccg	1380
ccgactcctc	cgtatgttta	tagggacacg	tgtaaactcg	gcctggccag	cctgagactg	1440
aaagcaaagc	agcactccag	cttcgggtac	gccagcgtgc	agaacccggc	ctccaacctg	1500
agtgccttgc	agtatgcagt	ggaccggccc	gtgtgagccg	caccacacag	gccgggatcc	1560
taggaccttg	ccggatgggg	caactccgcc	cttgaaagac	tgggaattat	gctagaaggt	1620
cgtgggcact	aaagaaaggg	agagaaagag	aagctatata	gagaaaagga	aaccactgaa	1680
tcaaagagag	agctcctttg	atttcaaagg	gatgtcctca	gtgtctgaca	tctttcacta	1740
caagtatttc	taacagttgc	aaggacacat	acacaaacaa	atgtttgact	ggatatgaca	1800
ttttaacatt	actataagct	tgttattttt	taagttagc	attgttaaca	tttaaataac	1860
tgaaaggatg	tatatatatc	gaaatgtcaa	attaatttta	taaaagcagt	tgtagtaaat	1920
atcacacacg	tgttttttaa	ggttaggctt	taaaataaag	catgtttata	agaagcgatt	1980
aggatttttc	gcttgcgagc	aaggagagtgt	atatactaaa	tgccacactg	tatgtttcta	2040
acatattatt	attattataa	aaaatgtgtg	aatatcagtt	ttagaatagt	ttctctgggtg	2100
gatgcaatga	tgtttctgaa	actgctatgt	acaacctacc	ctgtgtataa	catttcgtac	2160
aatattattg	ttttactttt	cagcaaatat	gaaacaaatg	tgttttattt	catgggagta	2220
aatatactg	catac					2235

<210> 21
<211>
<212> protein
<213> polypeptide Sequence

<220>
<223> amino acid sequence (Homo sapiens)

<400> 450

METNCRKLVSAACVQLGVQPAAVECLFSKDSEIKKVEFTDSPESRKEAASSKFFP
RQHPGANEKDKSQGKNEDVGAEDPSKKKRQRRQRTHTSQQQLQELEATFQRNR
YPDMSTREEIIVVWTLNTEARVRVWFKNRRRAKWRKRERNQQAELCKNGFGPQFNG
LMQPYDDMYPGYSYNWAAKGLTSASLSTKSFPPFNSMNVNPLSSQSMFSPNS
ISSMSMSSSMVPSAVTGVPSSLNSLNNLNNLSSPSLNSAVPTPACPYAPPTPP
YVYRDTCNSSLASRLKAKQHSSFGYASVQNPASNLSACQYAVDRPV

450

<210> 22
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)

<400> 22

gtaggggagg gaagtagatg t

21

<210> 23
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)

<400> 23

tcctcaactc tacaaccta aaa

23

<210> 24
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)

<400> 24
agtcgggaga gcgaaa 16

<210> 25
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)

<400> 25
agttgggaga gtgaaa 16

<210> 26
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)

<400> 26
aagagtcggg agtcgga 17

<210> 27
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)

<400> 27
aagagttggg agttgga 17

<210> 28
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)

<400> 28
ggtcgaagag tcggga 16

<210> 29
<211> 16

<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)

<400> 29

ggttgaagag ttggga

16

<210> 30
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)

<400> 30

atgttagcgg gtcgaa

16

<210> 31
<211> 16
<212> DNA
<213> Artificial Sequence

<220>
<223> chemically treated genomic DNA (Homo sapiens)

<400> 31

tagtgggttg aagagt

16

